

1-1-2020

Sleep duration is associated with vitamin D deficiency in older women living in Macao, China: A pilot cross-sectional study

Xiaoying Liu
University of Sydney

Liang Ke
Macao Hypertension League

Jacky Ho
Macao Hypertension League

Myriam Abboud
Zayed University, myriam.abboud@zu.ac.ae

Elias Mpofu
University of Sydney

See next page for additional authors

Follow this and additional works at: <https://zuscholars.zu.ac.ae/works>



Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Liu, Xiaoying; Ke, Liang; Ho, Jacky; Abboud, Myriam; Mpofu, Elias; Brennan-Speranza, Tara C.; Mason, Rebecca S.; and Brock, Kaye E., "Sleep duration is associated with vitamin D deficiency in older women living in Macao, China: A pilot cross-sectional study" (2020). *All Works*. 3103.
<https://zuscholars.zu.ac.ae/works/3103>

This Article is brought to you for free and open access by ZU Scholars. It has been accepted for inclusion in All Works by an authorized administrator of ZU Scholars. For more information, please contact scholars@zu.ac.ae.

Author First name, Last name, Institution

Xiaoying Liu, Liang Ke, Jacky Ho, Myriam Abboud, Elias Mpofu, Tara C. Brennan-Speranza, Rebecca S. Mason, and Kaye E. Brock

RESEARCH ARTICLE

Sleep duration is associated with vitamin D deficiency in older women living in Macao, China: A pilot cross-sectional study

Xiaoying Liu¹, Liang Ke², Jacky Ho^{2,3}, Myriam Abboud⁴, Elias Mpofu^{5,6,7}, Tara C. Brennan-Speranza^{1,8}, Rebecca S. Mason¹, Kaye E. Brock^{1*}

1 Department of Physiology, School of Medical Sciences, Bosch Institute, The University of Sydney, Sydney, Australia, **2** Macao Hypertension League, Macao, China, **3** City University of Macao, Macao SAR, China, **4** Zayed University, Dubai, UAE, **5** Clinical and Rehabilitation Sciences Research Group, Faculty of Health Science, The University of Sydney, Sydney, Australia, **6** Rehabilitation and Health Services, University of North Texas, Denton, TX, United States of America, **7** Educational Psychology, University of Johannesburg, Johannesburg, South Africa, **8** School of Public Health, Faculty of Medicine and Health, University of Sydney, Sydney, Australia

* Kaye.Brock@sydney.edu.au



OPEN ACCESS

Citation: Liu X, Ke L, Ho J, Abboud M, Mpofu E, Brennan-Speranza TC, et al. (2020) Sleep duration is associated with vitamin D deficiency in older women living in Macao, China: A pilot cross-sectional study. PLoS ONE 15(3): e0229642. <https://doi.org/10.1371/journal.pone.0229642>

Editor: Corrie Whisner, Arizona State University, UNITED STATES

Received: June 11, 2019

Accepted: February 11, 2020

Published: March 4, 2020

Copyright: © 2020 Liu et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Data Availability Statement: All relevant data are within the manuscript and its Supporting Information files.

Funding: Funding for this project was obtained from The Macao Hypertension League (2011–14) and The China Studies Center, University of Sydney. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing interests: The authors have declared that no competing interests exist.

Abstract

Chinese women are known to have both a high prevalence of metabolic syndrome (MetS) and vitamin D deficiency (serum 25-hydroxyvitamin D (25OHD) <50 nmol/l). Associations between sleep duration and circulating 25OHD have recently been reported but, to our knowledge, these associations have not been studied in older Chinese populations. We thus investigated whether sleep duration was associated with vitamin D status in a population from Macao, China, and whether sleep duration modified the association between MetS and vitamin D deficiency. In 207 older (>55 years) Macanese, anthropometry, blood samples and validated questionnaires, including sleep duration and cardiovascular risk factors, were simultaneously collected. On multivariable categorical analyses, those women, not men, who had short sleep duration (≤ 6 hours (h)) were at a 2-fold risk for vitamin D deficiency (both <50 nmol/L and <37 nmol/L; OR = 1.94, 95%CI 1.29–2.92; OR = 2.05, 95%CI 1.06–3.98, respectively) and those who had longer sleep duration (>8 h) were 3-fold more likely to have vitamin D deficiency (OR = 3.07, 95%CI 1.47–6.39; OR = 2.75, 95%CI 1.08–7.00, respectively) compared to those with normal sleep duration (6–8 h). Both women and men with MetS were 2-fold more likely to have vitamin D deficiency (women: OR = 2.04, 95%CI 1.31–3.17; OR = 2.15, 95%CI 1.11–4.17, respectively; men: OR = 2.01, 95%CI 1.23–3.28; OR = 2.04, 95%CI 1.00–4.29, respectively). Moreover, women with both short sleep duration and MetS had an increased risk of vitamin D deficiency (OR = 3.26, 95%CI 1.10–9.64). These associations were not found in those with longer sleep. Men with longer sleep and MetS had a 5-fold risk of vitamin D deficiency (OR = 5.22; 95%CI 2.70–10.12). This association was non-significant for men with shorter sleep. We conclude that both short and long sleep duration were associated with vitamin D deficiency in older Chinese women. Further research is needed in larger cohorts or with intervention studies to further examine the associations between reduced sleep, metabolic syndrome and vitamin D deficiency.

Introduction

Vitamin D is a hormone precursor, necessary for facilitating the absorption of calcium and phosphate ions from the gut [1]. Vitamin D deficiencies lead to osteomalacia, or rickets in children, both of which are problems of impaired bone mineralization. More recent studies have also indicated that vitamin D is beneficial for the prevention of skin cancer [2–4], and for improving muscle function and other physiological and pathophysiological processes [5–8]. In humans, following exposure to ultraviolet B radiation from the sun, 7-dehydrocholesterol is converted to Vitamin D. Vitamin D can also be obtained from the diet, however, skin production is more important in humans. Vitamin D status is defined by the amount of circulating 25-hydroxyvitamin D (25OHD). Vitamin D insufficiency (25OHD < 50 nmol/L) varies across populations worldwide [9–11] and is consistently reported to be common in Chinese populations [12–26], especially in women [12, 15, 21, 22, 24, 25, 27, 28]. Investigations into the factors associated with vitamin D deficiency in Chinese women have indicated that the traditional culture of using umbrellas and/or sunscreen to avoid tanning results in a lack of sunlight exposure leading to reduced vitamin D levels [12, 27].

In addition, a recent meta-analysis of epidemiological observational ($n = 849,412$) and randomized intervention studies ($n = 30,716$) has reported an inverse association of circulating 25OHD with risk of cardiovascular (CVD) death (Odds Ratio [OR] = 1.35, 95%CI 1.13–1.61) [29] and Chinese populations are reported to have an increasingly high prevalence of CVD [30]. Other data from this same study from Macao on associations between CVD risk and vitamin D deficiency has been previously published [31]. Cardiovascular mortality is also reported to be associated with both short and long sleep duration [32]. Three recent meta-analyses reported associations between CVD mortality and sleep duration. Both short sleep duration (Relative Risk [RR] = 1.18, 0.76–1.84) and long sleep duration (RR = 1.43, 1.15–1.78) were associated with CVD mortality [33–35]. In addition, metabolic syndrome, a cluster of risk factors for CVD (central obesity, reduced high-density lipoprotein cholesterol [HDL], increased blood pressure, blood glucose and blood triglycerides [TG]), were also reported to be associated with both short and long sleep duration [36, 37].

Vitamin D deficiency is associated with sleep apnoea [38] and sleep duration, however, there is conflicting data from intervention trials on the effect of vitamin D supplementation on sleep quality in men and women [39]. To our knowledge, no study has investigated the association between sleep duration and vitamin D status in adult Chinese populations. We have previously reported associations between vitamin D and CVD risk factors in a representative population from Macao, China ($n = 566$) [28]. As previously noted, metabolic syndrome is a risk factor for vitamin D deficiency [36, 37]. Very few studies have examined the interaction effect of short or long sleep duration and CVD risk factor on vitamin D deficiency. We thus took the opportunity to investigate sleep duration and vitamin D deficiency and CVD risk in those age >55 years from this population ($n = 207$) taking into account their metabolic syndrome status. We also examined whether sleep duration modified the association between metabolic syndrome and vitamin D deficiency. As older females had a much higher rate of vitamin D deficiency compared to men (47% in female vs. 26% in male) [12], the specific aims of this pilot were to investigate the relationship between vitamin D status and short (≤ 6 h) or long (> 8 h) sleep duration compared to normal sleep duration (6 to 8 h) [40]) in both men and women.

Materials and methods

Study sample and data collection

A representative population sample from Macao was investigated in 2014, detailed methods of sampling and data collection by individual measurement and validated questionnaires have

previously been documented [12, 28, 41]. In brief, based on an estimated hypertension prevalence of 28% in 2012, a sample size of 1,488 was predicted; with an estimated 50% response rate, 2,400 households (approximately 2.5 members per household) were randomly selected from a register of 170,000 households. A total of 1,410 study participants from 2,174 eligible participants were recruited (response rate 68%) and had completed measurement and survey. Of these 1,410 participants, 566 participants had donated blood samples for analyzing serum 25OHD and metabolic parameters. We had reported that the prevalence of vitamin D insufficiency was 55% in the total population, and it was estimated that the prevalence of vitamin D <50 nmol/L was 45% in China [42]. The post-hoc power was 99%. In these 566 participants, thirty-seven percent were older population ($n = 207$; age >55 years) (Fig 1). Furthermore, two older females and one male had no data on sleep duration, therefore, only 204 participants were included in the current analysis.

Sleep duration was documented as number of hours (h) of sleep per day on weekdays and weekends. We scaled the question responses retaining the intent of the Pittsburgh Sleep Questionnaire Index item [43] making for greater precision to guide our analysis. Daily average sleep duration was then calculated = [(sleep duration in weekday*5 + sleep duration in weekend*2) / 7]. A comparison of short sleep (≤ 6 h) and long sleep duration (>8 h) were compared to normal sleep duration (6–8 h). Although this normal duration differs slightly from the recommendations of the US National Sleep Foundation and American Academy of Sleep Medicine [44], the proposed 6–8 h “normal” sleep duration for this population comes from recent studies of the association between sleep duration and CVD events in Chinese population [45, 46].

Concurrently (in summer) fasting blood samples were collected and serum 25OHD was determined using an electrochemiluminescence immunoassay (Roche Diagnostics) with an intra-assay coefficient of variation of 1.93%. As opinion regarding optimal levels of vitamin D is divided [11, 47]. Chinese populations are known to have quite low mean values of vitamin D. Thus, vitamin D insufficiency was defined as serum 25OHD <50 nmol/L and vitamin D deficiency was defined as serum 25OHD <37 nmol/L in the current investigation.

Serum high-density lipoprotein (HDL) levels were determined using homogeneous enzymatic colorimetric assays and serum triglycerides (TG) were determined using an enzymatic colorimetric assay. Metabolic syndrome (MetS) was defined as the presence of any three of the following: overweight ($\text{BMI} \geq 24 \text{ kg/m}^2$), hypertension (measured blood pressure $>140/90$ mmHg and/or self-reported diagnosis and/or treatment of hypertension in the last two weeks), low serum HDL (serum HDL <1.03 mmol/L) or high TG (serum TG ≥ 1.7 mmol/L) (National Cholesterol Education Program Adult Treatment Panel III criteria) [48].

Socioeconomic status (SES), physical activity (PA), and fish consumption data were collected from individually administered questionnaires [12, 28, 41].

Socioeconomic status was adopted from area coding according to the income level of resident area [49].

Physical activity was measured by the International Physical Activity Questionnaire Short Version [50]. The walking, moderate and vigorous metabolic equivalent task (MET) minutes per week and the total MET minutes per week were calculated according to the IPAQ scoring protocol.

Information regarding oily fish consumption was collected by asking participants “Do you usually (≥ 4 times/week) eat fatty fish?”

Individual questionnaire survey and blood sample collection were conducted at the one visit. Ethics approval was obtained from both the Human Research Ethics Committee of the University of Sydney, Australia and Sun Yet-Sen University China. Informed consent was obtained in written form.




Fig 1. Flow diagram of study participants.

<https://doi.org/10.1371/journal.pone.0229642.g001>

Statistical analysis

The present sample was selected using a complex sample design where the study subjects were sampled with differential selection probabilities. Non-response and post-stratification adjustments were conducted to obtain the sample weights (the sample weights are the number of observations in the population that can be represented by the sampled observations) [51]. Survey data was weighted, and age and sex standardization was performed based on the population distribution from the Macao 2011 census, and thus, the selected sample was representative of the Macao population [41].

Data were presented as mean and standard deviation (SD) for continuous variables; and as proportions for categorical variables. Differences of population characteristics between older and younger population was calculated by chi-square test. Levels of serum 25OHD and sleep duration in those with short (≤ 6 h) or normal (>6 h and ≤ 8 h) or long (>8 h) sleep duration, and MetS were calculated using student *t*-tests. Vitamin D risk factors included SES, PA, and fish consumption [12, 28, 31] and MetS. In our conditional logistic regression model investigating vitamin D deficiency in this population, each risk factor variable (SES, PA and fish consumption) was added individually and if still significant, left in the model as a confounder.

The association between sleep duration and serum 25OHD levels was firstly assessed by quadratic regression stratified by sex. Quadratic term of sleep duration was added to the linear regression to test the quadratic association between sleep duration and serum 25OHD levels.

As an inverted 'u-shape' (or biphasic) relationship was observed between sleep duration and serum 25OHD levels, the associations between sleep and vitamin D deficiency were then assessed by univariable and multivariable logistic regression analyses. In detail, associations with vitamin D deficiency (as defined as both less than 37 nmol/L or less than 50 nmol/L) were then assessed by multivariable conditional logistic regression analyses, adjusting for SES, PA, and fish consumption and presented as odds ratios (OR) with 95% confidence intervals (CI). In this population from Macao, there were marked differences for predictors of vitamin D deficiency between the sexes (we previously reported significantly statistical interaction [12]); thus these data were stratified by sex.

In this current population, we also tested for multiplicative statistical interaction between sleep duration and MetS with vitamin D status. This was assessed by the likelihood ratio test in a logistic model [52] for vitamin D status, which included main effects for sleep duration (≤ 6 h) and MetS and the interaction term between sleep duration and MetS (Yes/No). As this was highly significant ($p < 0.001$), subsequent models were presented stratified by short (≤ 6 h) or normal-long (>6 h) sleep duration. According to our criteria, none of the older male participants with < 6 h sleep had vitamin D levels < 37 nmol/L fulfilled the criteria of MetS. Thus, analysis of this group was not possible, unless we classified one of the participants that was borderline for MetS as having MetS. As such, these results are based on the latter assumption.

For all analysis, statistical significance was defined as $p < 0.05$. All data analyses were performed with SPSS 21 statistical package (IBM, Armonk, NY)

Results

We have previously reported associations between vitamin D status and CVD risk factors in a representative population from Macao, China ($n = 566$) [28]. Table 1 shows the participant characteristic by age groups. Although the older population had lower prevalence of vitamin D insufficiency and severe deficiency compared to the younger population (36% vs. 63% and 10% vs. 28%, respectively), CVD risk factors, including overweight (48% vs. 34%), TG (30% vs. 20%), hypertension (61% vs. 20%) and MetS (21% vs. 9%) were higher in the older population.

Table 1. Characteristics of Macao population (n = 566) stratified by age.

	Older (n = 207)	Younger (n = 359)
25OHD <50 nmol/L	36	63*
25OHD <37 nmol/L	10	28*
Sleep <6h /day	32	19*
BMI \geq 24 kg/m ²	48	34*
Decreased HDL	20	17
Increased TG	30	20*
Hypertension	61	20*
MetS	21	9*
Mod-high PA	79	72*
Oily fish <4 time/wk	32	28*
Low SES	35	44*
Female	53	64*

Data were presented as %

*p<0.05 compared proportion between older and younger Macanese populations

<https://doi.org/10.1371/journal.pone.0229642.t001>

In addition, the rate of short sleep duration (<6 h/day) was significantly higher in the elderly than the younger (32% vs. 19%).

We thus investigated sleep duration and vitamin D status and CVD risk the older population from this population (n = 207). There were significant differences in vitamin D deficiency rates (serum 25OHD <50 nmol/L and <37 nmol/L) in women versus men in this population. The prevalence of serum 25OHD <50nmol/L was 47% in women vs. 26% in men (p<0.001). The prevalence of serum 25OHD <37 nmol/L was 12% in women vs. 8% in men (p<0.001). The mean serum 25OHD levels were 62.76 nmol/L for men and 54.46 nmol/L for women (p<0.001).

In older Macanese females, when the association between sleep duration and serum 25OHD levels was assessed by the quadratic regression, both sleep duration and the quadratic term were significantly associated with serum 25OHD levels ($y = 20.46 + 8.75x - 0.54x^2$, p<0.05 for both coefficients; $R^2 = 0.03$). As shown in Fig 2A, both long sleep duration and short sleep duration were associated with decreased serum 25OHD levels in these older Macanese women. In contrast, both sleep duration and the quadratic in older men were not associated with serum 25OHD levels ($y = 84.09 - 5.28x + 0.29x^2$; p>0.05 for both coefficients; $R^2 = 0.015$; Fig 2B).

In women, on conditional regression analysis, those with short and long sleep duration were at a 2- to 3-fold increased risk of vitamin D deficiency (either <50 nmol/L or <37 nmol/L) compared to those with normal sleep (short sleep duration OR = 1.94, 95% CI 1.29–2.92 and OR = 2.05, 95% CI 1.06–3.98 respectively, and longer sleep duration OR = 3.07, 95% CI 1.47–6.39 and OR = 2.75, 95% CI 1.08–7.00 respectively) (Table 2).

In men, shorter sleep (\leq 6 h) was not significantly associated with vitamin D deficiency (either <50 nmol/L or <37 nmol/L) in men. Longer sleep (>8h) was associated with serum 25OHD <37 nmol/L (OR = 3.01, 95%CI 1.22–7.42) but not associated with serum 25OHD <50 nmol/L (Table 2).

In addition, having MetS in both vitamin D deficiency categories was associated with 2-fold significant adjusted risk of vitamin D deficiency (either <50 nmol/L or <37 nmol/L) in both women and men (women: OR = 2.04, 95%CI 1.31–3.17; OR = 2.15, 95%CI 1.11–4.17, respectively) (men: OR = 2.08, 95%CI 1.23–3.20; OR = 2.04, 95%CI 1.00–4.29, respectively) (Table 2).




Fig 2. A. Sleep duration and serum 25OHD in older women living in Macao, China (n = 108). B. Sleep duration and serum 25OHD in older men living in Macao, China (n = 96).

<https://doi.org/10.1371/journal.pone.0229642.g002>

As a significant interaction between sleep duration and MetS was detected in both women and men ($p_{\text{interaction}} < 0.001$), these data were then stratified by short (≤ 6 h/day) or normal-long sleep duration (> 6 h/day) (Table 3).

On multivariable analysis, women with short sleep duration and having MetS had a significant 3-fold increased risk of vitamin D deficiency (serum 25OHD < 50 nmol/L and < 37 nmol/L) (OR = 3.26, 95%CI 1.10–9.64). The risk increased to 20-fold when severe vitamin D deficiency was investigated (OR = 37.53, 95%CI 7.06–199.61). It should be noted that only 35 women had short sleep duration, therefore, the small sample size led to the high OR and wide CI when we defined the cut-off for vitamin D deficiency as < 37 nmol/L. In contrast, these associations were not found in those women with longer sleep. (Table 3).

The opposite effect was seen in men, where on multivariable analysis those with longer sleep and MetS had a 5-fold risk of vitamin D deficiency (serum 25OHD < 37 nmol/L, OR = 4.86; 95%CI 2.09–11.33; serum 25OHD < 50 nmol/L, OR = 5.22; 95%CI 2.70–10.12) and a non-significant association between MetS and vitamin D deficiency for those with shorter sleep.

Discussion

The current investigation of an older Macanese population found that both short and long sleep duration were associated with increased risk of vitamin D deficiency in women, but not in men.

Although there have been some clinical evidence linking sleep apnoea and vitamin D deficiency [38], few studies have investigated the association between sleep duration and circulating 25OHD levels or dietary vitamin D intake [53–66] (Table 4).

Of the two studies [55, 57] that investigated vitamin D supplementation and change in sleep duration, one small study (n = 28) reported that chronic pain patients with serum 25OHD < 50 nmol/L had a significantly shorter sleep duration than those with serum 25OHD between 50–75 nmol/L at baseline (3.73 hours vs. 5.33 hours); these chronic pain patients reported increased sleep duration after three months of vitamin D supplementation [55]. It should be noted, however, that our study had a higher average sleep duration than study by Huang *et al.*, due to the difference in participant characteristics. The findings regarding the association between shorter sleep duration and increased risk of vitamin D deficiency were consistent. In contrast, the authors of the randomized control trail undertaken in overweight,

Table 2. Sleep duration (h) and risk of serum 25OHD <50 nmol/L or 25OHD <37 nmol/L in older Macao residents (n = 204).

	%	Mean sleep duration (h)	Mean 25OHD (nmol/L)	25OHD <50 nmol/L		25OHD <37 nmol/L	
				Crude OR (95% CI)	Adjusted OR ^a (95% CI)	Crude OR (95% CI)	Adjusted OR ^b (95% CI)
Women (n = 108)							
Sleep duration (h)							
Sleep >6 to ≤8 (reference)	61	7.5 ± 0.5	56 ± 17	1.0	1.0	1.0	1.0
Sleep ≤6	30	5.3 ± 1.0*	52 ± 17*	1.55 (1.05–2.30)*	1.94 (1.29–2.92)*	1.47 (0.80–2.70)	2.05 (1.06–3.98)*
Sleep >8	8	9.2 ± 0.5*	50 ± 23	2.25 (1.12–4.54)*	3.07 (1.47–6.39)*	2.89 (1.24–6.74)*	2.75 (1.08–7.00)*
MetS							
No (reference)	75	6.9 ± 1.3	56 ± 18	1.0	1.0	1.0	1.0
Yes	25	7.2 ± 1.3	50 ± 16*	1.95 (1.28–2.95)*	2.04 (1.31–3.17)*	1.55 (0.85–2.82)	2.15 (1.11–4.17)*
Men (n = 96)							
Sleep duration (h)							
Sleep >6 to ≤8 (reference)	55	7.4 ± 0.6	60 ± 15	1.0	1.0	1.0	1.0
Sleep ≤6	31	5.4 ± 1.0*	65 ± 19	0.89 (0.57–1.39)	0.75 (0.48–1.20)	0.98 (0.46–2.10)	0.85 (0.39–1.84)
Sleep >8	14	9.5 ± 0.7*	65 ± 23	0.45 (0.21–1.00)	0.56 (0.26–1.21)	2.28 (0.98–5.31)	3.01 (1.22–7.42)*
MetS							
No (reference)	80	7.2 ± 1.5	65 ± 18	1.0	1.0	1.0	1.0
Yes	20	6.5 ± 1.4*	56 ± 15*	2.30 (1.45–3.65)*	2.01 (1.23–3.28)*	1.99 (0.99–3.99)	2.04 (1.00–4.29)*

*p<0.05 for independent t-test and logistic regression compared to the reference group

^a Mutually adjusted for sleep duration, MetS, SES, PA, fish consumption

^b Mutually adjusted for sleep duration, MetS, SES and PA; Data presented as mean +SD where appropriate

<https://doi.org/10.1371/journal.pone.0229642.t002>

vitamin D deficient postmenopausal women, did not observe improvements in sleep duration and sleep quality after 12 months of vitamin D supplementation [57].

Of the eleven observational studies investigating sleep duration, [53, 54, 56, 58–65], measured either by polysomnography [64], actigraphy [53, 58, 61] or as self-reported [54, 56, 59, 60, 62, 63], six large epidemiological cross-sectional studies reported a significant crude association between short sleep duration and decreased serum 25OHD levels [53, 54, 56, 58, 62, 64]. Three of these studies were from the USA and appeared to have slightly shorter average sleep duration and a higher mean 25OHD levels than our study [53, 54, 58]. Bertisch *et al.* reported a mean serum 25OHD level at 63 nmol/L and a mean sleep duration of 6.5 h/day [53]; Beydoun *et al.*, reported a mean serum 25OHD level at 55 nmol/L and 37% had short sleep duration [54]; Massa *et al.*, reported only 16% of the participants had vitamin D deficiency and the mean sleep duration was 6.4 h/day [58]. However, three Korean studies had similar sleep duration (mean sleep durations were 6.6, 7.1 and 6.6 h/day, respectively) and slightly lower mean 25OHD levels (all Korean studies reported mean serum 25OHD levels <50 nmol/L), although it should be noted that the population were older than in our study (all age >65 years) [56, 59, 62]. To our knowledge, study of the association between sleep duration and vitamin D status has been limited in Chinese population. A recently published paper from Ningbo, China, reported that vitamin D deficiency was associated with a two-fold likelihood of short sleep duration (< 9h) in school children [65]. In addition, a recent meta-analysis [66] reported that having low serum vitamin D levels was associated with increased risk of short sleep duration (pool estimate OR = 1.74; 95%CI 1.30–2.23). None of the included studies were in Chinese populations [66].

Table 3. Serum 25OHD <50 nmol/L or serum 25OHD <37 nmol/L risk stratified by duration of sleep (≤ 6 h and >6 h) in older Macanese (n = 204).

	%	Mean 25OHD (nmol/L)	25OHD <50 nmol/L		25OHD <37 nmol/L	
			Crude OR (95%CI)	Adjusted OR (95% CI)	Crude OR (95%CI)	Adjusted OR (95% CI)
Women (n = 108)						
Sleep ≤ 6 h (n = 35)						
MetS						
No (reference)	79	54 \pm 17	1.0	1.0	1.0	1.0
Yes	21	42 \pm 13*	3.34 (1.30–8.61)*	3.26 (1.10–9.64)*	5.36 (1.96–14.71)*	37.53 (7.06–199.61)*
Sleep >6 h (n = 73)						
MetS						
No (reference)	72	57 \pm 19	1.0	1.0	1.0	1.0
Yes	28	53 \pm 16	1.76 (1.09–2.84)*	1.41 (0.85–2.35)	0.83 (0.37–1.84)	0.64 (0.27–1.53)
Men^b (n = 96)						
Sleep ≤ 6 h (n = 30)						
MetS						
No (reference)	68	65 \pm 22	1.0	1.0	1.0	1.0
Yes	32	64 \pm 11	0.49 (0.22–1.12)	0.49 (0.20–1.21)	1.99 (0.60–6.57)	2.47 (0.69–8.80)
Sleep >6 h (n = 66)						
MetS						
No (reference)	84	63 \pm 16	1.0	1.0	1.0	1.0
Yes	16	48 \pm 14*	6.62 (3.49–12.56)*	5.22 (2.70–10.12)*	5.93 (2.63–13.35)*	4.86 (2.09–11.33)*

*p<0.05 for independent t-test and logistic regression compared to the reference group

^a Mutually adjusted for MetS, SES, PA and fish consumption

^b Mutually adjusted for MetS, SES, and PA; Data presented as mean+ SD where appropriate

<https://doi.org/10.1371/journal.pone.0229642.t003>

Of the three studies that investigated longer sleep duration [54, 56, 60], only one study (n = 93) reported an inverse correlation between sleep duration and serum 25OHD status ($r = -0.0261$) [60]. In addition, when investigating intake of dietary vitamin D, one large epidemiological US National Health and Nutritional Study reported significant lower dietary vitamin D intake mean associations between both long and short sleep [63].

Circadian entrainment research has indicated that the sleep-wake cycle is reset daily by retina-perceived light, controlled by a master clock in the suprachiasmatic nuclei of the hypothalamus that inhibits the release of melatonin [67]. One uncontrolled clinical trial reported that vitamin D supplementation in 1,500 patients with sleep disturbances resulted in normal sleep [68]. These authors thus linked sleep disturbances with vitamin D deficiency and suggested that 25OHD deficiency may be involved in circadian rhythm dysynchronization [68].

Metabolic syndrome has been reported to be associated with vitamin D status in a recent meta-analysis (Top vs. bottom tertile vitamin D levels, RR = 0.86, 95%CI 0.80–0.92; n = 6554) [69]. To our knowledge, this is the first time MetS has been associated with sleep duration (either short or long) and vitamin D deficiency. However, recently published data from an elderly Korean population showed a higher risk for obesity in those who were vitamin D deficient and had shorter sleep duration [62] (Table 4).

It is interesting that in our data, women with shorter sleep had a higher risk of vitamin D deficiency if MetS was present but men with MetS and longer sleep had a higher risk of vitamin D deficiency. We believe that this finding may be explained by the very high rate of vitamin D deficiency in women compared to men in this study population.

One explanation for the observations in the current study may be found in recent research that indicates the long duration of residence for the main vitamin D metabolite, 25(OH)D, in

Table 4. Studies investigating the association between sleep duration and vitamin D (serum 25OHD levels or dietary intake).

Author, year	N	%F	Country/ Ethnicity & Study design	Sleep duration (mean ± SD)/ methods	25OHD (nmol/L) (mean ± SD or median [IQR])	Age (years) (mean / range)	Univariable association between sleep duration and 25OHD or dietary VD intake
Mason, 2016 [57]	218	100	USA; RCT	PSQI sub-score for sleep duration;	53 ± 15	60 ± 5	No difference in change in sleep duration between placebo and VD suppl. group; ↔
Huang et al., 2013 [55]	28	64	USA; IN	4.5 ± 1.5; SR	46 ± 14	46 ± 11	↑ sleep duration after VD suppl. for 3 months;
Bertisch, et al., 2015 [53]	1721	55	USA; CS	6.5 ± 1.3; wrist actigraphy device;	63 ± 26	68 ± 9	↓ sleep duration in ↓ 25OHD < 50 nmol/ L participants vs. 25OHD > 75 nmol/L
Massa et al., 2015 [58]	2966	0	USA; CS	6.4 ± 1.2 h; wrist actigraphy device;	16% 25OHD < 50;	> 68	↓ sleep duration as ↓ 25OHD;
Piovezan, et al., 2017 [64]	657	56	Brazil; CS	46% had < 6h sleep; polysomnographic;	60% 25OHD < 75;	52 ± 9	↓ sleep duration ↓ 25OHD;
Darling et al., 2018 [61]	41	100	UK / CS	Sleep duration NA; Actigraphy & SR	South Asian 53 [51]; White Caucasian 83 [36];	39–75	No association between sleep duration and 25OHD; ↔
Beydoun, et al., 2014 [54]	2459	NA	USA; CS	37% had < 6h sleep; SR	55 ± 3	20–85	↓ sleep duration ↓ 25OHD;
Kim, 2014 [56]	1614	54	Korea; CS	6.6 ± 1.6; 47% had < 6h sleep; SR	49 ± 19	68 ± 5	↓ sleep duration ↓ 25OHD;
Song, 2016 [59]	2853	66	Korea; CS	7.1 ± 1.5 h/day; SR	42 [30–56]	72 ± 5	↓ sleep duration ↓ 25OHD in men
Darling et al., 2011 [60]	90	100	UK; CS	Sleep duration NA; PSQI	NA	NA	↓ sleep duration ↑ 25OHD;
Doo M., 2018 [62]	3757	56	Korea; CS	6.6 ± 0.05; SR	48 [range 10–134]	65–97	↓ sleep duration ↓ 25OHD;
Gong, et al., 2018 [65]	800	46	China, CS	9.17 ± 0.97; SR	56 ± 15	11 ± 2	↓ sleep duration ↓ 25OHD;
Grandner et al., [63]	4548	53	USA; CS	Very short (< 5); Short (5–6); Normal (7–8); Long (≥ 9); SR	4.4 ± 5.0 mcg;	46 ± 17	↓ dietary VD intake in very short, short and long sleep duration compared to normal

Abbreviation: N, number of participants; F, female; 25OHD, 25-hydroxyvitamin D; SD, standard deviation; IQR, interquartile range; CS, cross-sectional; RCT, randomized control trial; IN, intervention; PSQI, Pittsburgh Sleep Quality Index; VD, vitamin D; suppl., supplementation; NA, not available; ↔, no association; ↑, increase; ↓, decrease; SR, self-report; Data presented as mean + SD where appropriate

<https://doi.org/10.1371/journal.pone.0229642.t004>

the circulation, is that it is taken up into muscle cells where it binds to vitamin D-binding protein (DBP) that is attached to cytoplasmic actin [70]. Thus, muscle acts as a storage center for 25(OH)D, where it is able to passage into and out of the muscle cells. There is evidence that this process is particularly useful for maintaining vitamin D status in the winter months when UV exposure is reduced. Furthermore, it has been proposed that this mechanism can be negatively affected when muscle function is compromised by inactivity or malnutrition. Sleep duration is well known to impact muscle function [71–73]. Taken together, it is likely that sleep requirements for adequate muscle function also affect the storage of vitamin D and if, over long periods of time, a person has disturbed sleep, a risk of vitamin D deficiency develops. Of note, the significant differences we observed here in the 25(OH)D mean values suggest physiological changes in the mechanisms controlling blood 25(OH)D levels, rather than indications of relative sufficiency or deficiency of functional vitamin D. Indeed, one study in rats has shown that the relative levels of muscle-active hormones is related to sleep [74]. Thus, a future follow-up study to further enhance the understanding of the mechanism behind our current findings would be to measure various blood hormone levels in participants with sleep

deprivation and determine any correlations in these endocrine changes to changes in muscle handling of 25(OH)D.

The limitations of the current study include the cross-sectional study design and small sample size (especially when stratified). As with all cross-sectional studies, it is impossible to disentangle causality. Central obesity and fasting blood glucose were not able to be included in our definition of metabolic syndrome, as these measurements were not taken at the study visits. To address this, a BMI cut off at 24 kg/m² was applied in the current study. Several studies from China have suggested the optimal cut-off points to predict the presence of metabolic syndrome in adults were BMI between 23–24 kg/m² [75–78]. In addition, information on other sleep-related disturbances or disorders, including sleep apnoea, were not collected in our study. Several studies have reported that sleep apnoea was associated with lower serum 25OHD levels [79–87]. Furthermore, it remains possible that the associations between sleep and vitamin D status may be due to associations with poor health [88], including unmeasured confounding variables such as depression [89] and musculoskeletal pain [39] or inflammation [90]; these unmeasured confounding variables may differ by sex especially in Chinese populations. The strengths of this investigation are the representativeness of the population and that many risk factors for CVD were individually measured, enabling us to calculate MetS.

Conclusion

Both short and long sleep duration were associated with vitamin D deficiency in the older Macao women. Notably, women with shorter sleep and MetS had a very high risk of vitamin D deficiency. While the underlying unmeasured factors determining sleep duration may also contribute to MetS risk and vitamin D levels, further studies especially stratified by sex in Asian populations may provide insight into the causality of these associations.

Supporting information

S1 Data.
(XLSX)

Acknowledgments

We would like to acknowledge J Feng, F Van, S Leong, W Lau, C Kowk and all participants, study interviewers and study assistants (School of Nursing) and Kiang Hu Hospital Department of Laboratory. We also acknowledged Prof. David R. Fraser for his insightful input and proofreading of this manuscript.

Author Contributions

Conceptualization: Liang Ke, Jacky Ho, Elias Mpofu, Rebecca S. Mason, Kaye E. Brock.

Data curation: Liang Ke, Jacky Ho.

Formal analysis: Xiaoying Liu, Liang Ke, Kaye E. Brock.

Funding acquisition: Liang Ke.

Investigation: Xiaoying Liu, Kaye E. Brock.

Methodology: Kaye E. Brock.

Supervision: Liang Ke, Myriam Abboud, Elias Mpofu, Tara C. Brennan-Speranza, Rebecca S. Mason, Kaye E. Brock.

Writing – original draft: Xiaoying Liu, Liang Ke, Jacky Ho, Myriam Abboud, Elias Mpofu, Tara C. Brennan-Speranza, Rebecca S. Mason, Kaye E. Brock.

Writing – review & editing: Xiaoying Liu, Liang Ke, Jacky Ho, Myriam Abboud, Elias Mpofu, Tara C. Brennan-Speranza, Rebecca S. Mason, Kaye E. Brock.

References

1. Bouillon R, Marcocci C, Carmeliet G, Bikle D, White JH, Dawson-Hughes B, et al. Skeletal and extra-skeletal actions of vitamin D: Current evidence and outstanding questions. *Endocrine Reviews*. 2018; er.2018-00126. <http://dx.doi.org/10.1210/er.2018-00126>.
2. Dixon KM, Deo SS, Wong G, Slater M, Norman AW, Bishop JE, et al. Skin cancer prevention: A possible role of 1,25dihydroxyvitamin D3 and its analogs. *The Journal of Steroid Biochemistry and Molecular Biology*. 2005; 97(1):137–43. <https://doi.org/10.1016/j.jsbmb.2005.06.006>.
3. Saw RPM, Armstrong BK, Mason RS, Morton RL, Shannon KF, Spillane AJ, et al. Adjuvant therapy with high dose vitamin D following primary treatment of melanoma at high risk of recurrence: a placebo controlled randomised phase II trial (ANZMTG 02.09 Mel-D). *BMC Cancer*. 2014; 14:780. <https://doi.org/10.1186/1471-2407-14-780> PMC4221702. PMID: 25343963
4. Rebecca SM, Jorg R. Sunlight Vitamin D and Skin Cancer. *Anti-Cancer Agents in Medicinal Chemistry*. 2013; 13(1):83–97. <http://dx.doi.org/10.2174/1871520611307010083>. PMID: 23094924
5. Meier C, Woitge HW, Witte K, Lemmer B, Seibel MJ. Supplementation With Oral Vitamin D3 and Calcium During Winter Prevents Seasonal Bone Loss: A Randomized Controlled Open-Label Prospective Trial. *Journal of Bone and Mineral Research*. 2004; 19(8):1221–30. <https://doi.org/10.1359/JBMR.040511> PMID: 15231008
6. Tanner SB, Harwell SA. More than healthy bones: a review of vitamin D in muscle health. *Therapeutic Advances in Musculoskeletal Disease*. 2015; 7(4):152–9. <https://doi.org/10.1177/1759720X15588521> PMC4530385. PMID: 26288665
7. Brennan-Speranza TC, Mor D, Mason RS, Bartlett JR, Duque G, Levinger I, et al. Skeletal muscle vitamin D in patients with end stage osteoarthritis of the knee. *The Journal of Steroid Biochemistry and Molecular Biology*. 2017; 173:180–4. <https://doi.org/10.1016/j.jsbmb.2017.01.022> PMID: 28161531
8. Mason C, Tapsoba JDD, Duggan C, Imayama I, Wang C-Y, Korde L, et al. Effects of vitamin D(3) supplementation on lean mass, muscle strength and bone mineral density during weight loss: A double-blind randomized controlled trial. *Journal of the American Geriatrics Society*. 2016; 64(4):769–78. <https://doi.org/10.1111/jgs.14049> PMC4840082. PMID: 27060050
9. Holick MF, Chen TC. Vitamin D deficiency: a worldwide problem with health consequences. *The American Journal of Clinical Nutrition*. 2008; 87(4):1080S–6S. <https://doi.org/10.1093/ajcn/87.4.1080S> PMID: 18400738
10. Palacios C, Gonzalez L. Is vitamin D deficiency a major global public health problem? *The Journal of Steroid Biochemistry and Molecular Biology*. 2014; 144:138–45. <https://doi.org/10.1016/j.jsbmb.2013.11.003> PMID: 24239505
11. Mithal A, Wahl DA, Bonjour J-P, Burckhardt P, Dawson-Hughes B, Eisman JA, et al. Global vitamin D status and determinants of hypovitaminosis D. *Osteoporosis International*. 2009; 20(11):1807–20. <https://doi.org/10.1007/s00198-009-0954-6> PMID: 19543765
12. Ke L, Mason RS, Mpofu E, Dibley M, Li Y, Brock KE. Vitamin D and parathyroid hormone status in a representative population living in Macau, China. *The Journal of Steroid Biochemistry and Molecular Biology*. 2015; 148:261–8. <https://doi.org/10.1016/j.jsbmb.2015.01.019> PMID: 25636721
13. Dorjgochoo T, Ou Shu X, Xiang Y-B, Yang G, Cai Q, Li H, et al. Circulating 25-hydroxyvitamin D levels in relation to blood pressure parameters and hypertension in the Shanghai Women's and Men's Health Studies. *British Journal of Nutrition*. 2012; 108(3):449–58. Epub 02/27. <https://doi.org/10.1017/S0007114511005745> PMID: 22365135
14. Lu H-K, Zhang Z, Ke Y-H, He J-W, Fu W-Z, Zhang C-Q, et al. High Prevalence of Vitamin D Insufficiency in China: Relationship with the Levels of Parathyroid Hormone and Markers of Bone Turnover. *PLOS ONE*. 2012; 7(11):e47264. <https://doi.org/10.1371/journal.pone.0047264> PMID: 23144810
15. Brock KE, Ke L, Tseng M, Clemson L, Koo FK, Jang H, et al. Vitamin D status is associated with sun exposure, vitamin D and calcium intake, acculturation and attitudes in immigrant East Asian women living in Sydney. *The Journal of Steroid Biochemistry and Molecular Biology*. 2013; 136:214–7. <https://doi.org/10.1016/j.jsbmb.2012.12.005> PMID: 23262263
16. Chen WR, Chen YD, Shi Y, Yin DW, Wang H, Sha Y. Vitamin D, parathyroid hormone and risk factors for coronary artery disease in an elderly Chinese population. *Journal of Cardiovascular Medicine*. 2015;

- 16(1):59–68. <https://doi.org/10.2459/JCM.0000000000000094> 01244665-201501000-00008. PMID: 24842466
17. Li L, Yin X, Yao C, Zhu X, Wu X. Vitamin D, Parathyroid Hormone and Their Associations with Hypertension in a Chinese Population. *PLOS ONE*. 2012; 7(8):e43344. <https://doi.org/10.1371/journal.pone.0043344> PMID: 22937036
 18. Lin S-W, Chen W, Fan J-H, Dawsey SM, Taylor PR, Qiao Y-L, et al. Prospective Study of Serum 25-Hydroxyvitamin D Concentration and Mortality in a Chinese Population. *American Journal of Epidemiology*. 2012; 176(11):1043–50. <https://doi.org/10.1093/aje/kws285> PMID: 23139250
 19. Lu L, Yu Z, Pan A, Hu FB, Franco OH, Li H, et al. Plasma 25-Hydroxyvitamin D Concentration and Metabolic Syndrome Among Middle-Aged and Elderly Chinese Individuals. *Diabetes Care*. 2009; 32(7):1278–83. <https://doi.org/10.2337/dc09-0209> PMID: 19366976
 20. Qiao Z, Li-xing S, Nian-chun P, Shu-jing X, Miao Z, Hong L, et al. Serum 25(OH)D Level and Parathyroid Hormone in Chinese Adult Population: A Cross-Sectional Study in Guiyang Urban Community from Southeast of China. *International Journal of Endocrinology*. 2013; 2013:150461. <https://doi.org/10.1155/2013/150461> PMC3771443. PMID: 24065989
 21. Woo J, Lam CWK, Leung J, Lau WY, Lau E, Ling X, et al. Very high rates of vitamin D insufficiency in women of child-bearing age living in Beijing and Hong Kong. *British Journal of Nutrition*. 2008; 99(6):1330–4. Epub 06/01. <https://doi.org/10.1017/S0007114507844382> PMID: 17961293
 22. Yan L, Prentice A, Zhang H, Wang X, Stirling DM, Golden MM. Vitamin D status and parathyroid hormone concentrations in Chinese women and men from north-east of the People's Republic of China. *European Journal Of Clinical Nutrition*. 2000; 54(1):68–72. <https://doi.org/10.1038/sj.ejcn.1600895> PMID: 10694775
 23. Yin X, Sun Q, Zhang X, Lu Y, Sun C, Cui Y, et al. Serum 25(OH)D is inversely associated with metabolic syndrome risk profile among urban middle-aged Chinese population. *Nutrition Journal*. 2012; 11(1):68. <https://doi.org/10.1186/1475-2891-11-68> PMID: 22958612
 24. Fang F, Wei H, Wang K, Tan L, Zhang W, Ding L, et al. High prevalence of vitamin D deficiency and influencing factors among urban and rural residents in Tianjin, China. *Archives of Osteoporosis*. 2018; 13(1):64. <https://doi.org/10.1007/s11657-018-0479-8> PMID: 29860553
 25. Zhen D, Liu L, Guan C, Zhao N, Tang X. High prevalence of vitamin D deficiency among middle-aged and elderly individuals in northwestern China: Its relationship to osteoporosis and lifestyle factors. *Bone*. 2015; 71:1–6. <https://doi.org/10.1016/j.bone.2014.09.024> PMID: 25284157
 26. Zhu W, Heil DP. Associations of vitamin D status with markers of metabolic health: A community-based study in Shanghai, China. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2018; 12(5):727–32. <https://doi.org/10.1016/j.dsx.2018.04.010>.
 27. Jang H, Koo F, Oo L, Clemson L, Cant R, Fraser D, et al. Culture and Sun Exposure in Immigrant East Asian Women Living in Australia 2013. 504–18 p.
 28. Ke L, Mason RS, Mpofu E, Vingren JL, Li Y, Graubard BI, et al. Hypertension and other cardiovascular risk factors are associated with vitamin D deficiency in an urban Chinese population: A short report. *The Journal of Steroid Biochemistry and Molecular Biology*. 2017; 173:286–91. <https://doi.org/10.1016/j.jsbmb.2016.11.011> PMID: 27865973
 29. Chowdhury R, Kunutsor S, Vitezova A, Oliver-Williams C, Chowdhury S, Kieft-de-Jong JC, et al. Vitamin D and risk of cause specific death: systematic review and meta-analysis of observational cohort and randomised intervention studies. *British Medical Journal*. 2014; 348:g1903. <https://doi.org/10.1136/bmj.g1903> PMID: 24690623
 30. Yang ZJ, Liu J, Ge JP, Chen L, Zhao ZG, Yang WY. Prevalence of cardiovascular disease risk factor in the Chinese population: the 2007–2008 China National Diabetes and Metabolic Disorders Study. *European Heart Journal*. 2012; 33(2):213–20. <https://doi.org/10.1093/eurheartj/ehr205> PMID: 21719451
 31. Ke L, Ho J, Feng J, Mpofu E, Dibley MJ, Feng X, et al. Modifiable risk factors including sunlight exposure and fish consumption are associated with risk of hypertension in a large representative population from Macau. *The Journal of Steroid Biochemistry and Molecular Biology*. 2014; 144:152–5. <https://doi.org/10.1016/j.jsbmb.2013.10.019> PMID: 24189545
 32. Full KM, Kerr J, Song D, Malhotra A, Gallo L, Arredondo EM, et al. Abstract 20971: Associations Between Objectively Measured Sleep Duration and 10-Year Predicted Cardiovascular Risk in Older Adult Women. *Circulation*. 2017; 136(suppl_1):A20971. https://doi.org/10.1161/circ.136.suppl_1.20971
 33. Silva AA, Mello RGB, Schaan CW, Fuchs FD, Redline S, Fuchs SC. Sleep duration and mortality in the elderly: a systematic review with meta-analysis. *BMJ Open*. 2016; 6(2):e008119. <https://doi.org/10.1136/bmjopen-2015-008119> PMID: 26888725
 34. Jike M, Itani O, Watanabe N, Buysse DJ, Kaneita Y. Long sleep duration and health outcomes: A systematic review, meta-analysis and meta-regression. *Sleep Medicine Reviews*. 2018; 39:25–36. <https://doi.org/10.1016/j.smr.2017.06.011> PMID: 28890167

35. Cappuccio FP, Cooper D, D'Elia L, Strazzullo P, Miller MA. Sleep duration predicts cardiovascular outcomes: a systematic review and meta-analysis of prospective studies. *European Heart Journal*. 2011; 32(12):1484–92. <https://doi.org/10.1093/eurheartj/ehr007> PMID: 21300732
36. Xi B, He D, Zhang M, Xue J, Zhou D. Short sleep duration predicts risk of metabolic syndrome: a systematic review and meta-analysis. *Sleep medicine reviews*. 2014; 18(4):293–7. <https://doi.org/10.1016/j.smrv.2013.06.001> PMID: 23890470
37. Hall MH, Muldoon MF, Jennings JR, Buysse DJ, Flory JD, Manuck SB. Self-Reported Sleep Duration is Associated with the Metabolic Syndrome in Midlife Adults. *Sleep*. 2008; 31(5):635–43. <https://doi.org/10.1093/sleep/31.5.635> PMID: 18517034
38. Neighbors CLP, Noller MW, Song SA, Zaghi S, Neighbors J, Feldman D, et al. Vitamin D and obstructive sleep apnea: a systematic review and meta-analysis. *Sleep Medicine*. 2018; 43:100–8. <https://doi.org/10.1016/j.sleep.2017.10.016> PMID: 29482804
39. de Oliveira DL, Hirotsu C, Tufik S, Andersen ML. The interfaces between vitamin D, sleep and pain. *Journal of Endocrinology*. 2017; 234(1):R23–R36. <https://doi.org/10.1530/JOE-16-0514> PMID: 28536294
40. Stone KL, Ewing SK, Ancoli-Israel S, Ensrud KE, Redline S, Bauer DC, et al. Self-Reported Sleep and Nap Habits and Risk of Mortality in a Large Cohort of Older Women. *Journal of the American Geriatrics Society*. 2009; 57(4):604–11. <https://doi.org/10.1111/j.1532-5415.2008.02171.x> PMID: 19220560
41. Ke L, Ho J, Feng J, Mpofo E, Dibley MJ, Li Y, et al. Prevalence, Awareness, Treatment and Control of Hypertension in Macau: Results From a Cross-Sectional Epidemiological Study in Macau, China. *American Journal of Hypertension*. 2015; 28(2):159–65. <https://doi.org/10.1093/ajh/hpu121> PMID: 25063734
42. Yu S, Fang H, Han J, Cheng X, Xia L, Li S, et al. The high prevalence of hypovitaminosis D in China: a multicenter vitamin D status survey. *Medicine*. 2015; 94(8):e585–e. <https://doi.org/10.1097/MD.0000000000000585> PMID: 25715263.
43. Buysse DJ, Reynolds CF III, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry research*. 1989; 28(2):193–213. [https://doi.org/10.1016/0165-1781\(89\)90047-4](https://doi.org/10.1016/0165-1781(89)90047-4) PMID: 2748771
44. Hirshkowitz M, Whiton K, Albert SM, Alessi C, Bruni O, DonCarlos L, et al. National Sleep Foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health*. 2015; 1(1):40–3. <https://doi.org/10.1016/j.sleh.2014.12.010> PMID: 29073412
45. Hu L, Huang X, Zhou W, You C, Li J, Li P, et al. Effect of hypertension status on the association between sleep duration and stroke among middle-aged and elderly population. *The Journal of Clinical Hypertension*. 2020; 22(1):65–73. <https://doi.org/10.1111/jch.13756> PMID: 31816157
46. Hu L, Zhang B, Zhou W, Huang X, You C, Li J, et al. Sleep duration on workdays or nonworkdays and cardiac–cerebral vascular diseases in Southern China. *Sleep Medicine*. 2018; 47:36–43. <https://doi.org/10.1016/j.sleep.2017.11.1147> PMID: 29880146
47. Dawson-Hughes B, Heaney RP, Holick MF, Lips P, Meunier PJ, Vieth R. Estimates of optimal vitamin D status. *Osteoporosis International*. 2005; 16(7):713–6. <https://doi.org/10.1007/s00198-005-1867-7> PMID: 15776217
48. Grundy SM, Cleeman JI, Daniels SR, Donato KA, Eckel RH, Franklin BA, et al. Diagnosis and Management of the Metabolic Syndrome. *Circulation*. 2005; 112(17):2735–52. <https://doi.org/10.1161/CIRCULATIONAHA.105.169404> PMID: 16157765
49. Government of Macao Special Administrative Region, Statistics and Census Service, Macao Yearbook of Statistics 2013 [Internet]. 2013. Available from: <https://www.dsec.gov.mo/en-US/Home/Publication/YearbookOfStatistics>.
50. Craig CL, Marshall AL, Sjoström M, Bauman AE, Booth ML, Ainsworth BE, et al. International physical activity questionnaire: 12-country reliability and validity. *Medicine and science in sports and exercise*. 2003; 35(8):1381–95. <https://doi.org/10.1249/01.MSS.0000078924.61453.FB> PMID: 12900694
51. Korn EL, Graubard BI. *Analysis of Health Surveys*: Wiley; 2011.
52. Breslow NE, Day NE, Schlesselman JJ. Statistical Methods in Cancer Research. Volume 1—The Analysis of Case-Control Studies. *Journal of Occupational and Environmental Medicine*. 1982; 24(4):255–7. 00005122-198204000-00006.
53. Bertisch SM, Sillau S, de Boer IH, Szklo M, Redline S. 25-Hydroxyvitamin D Concentration and Sleep Duration and Continuity: Multi-Ethnic Study of Atherosclerosis. *Sleep*. 2015; 38(8):1305–11. <https://doi.org/10.5665/sleep.4914> PMID: 25669179
54. Beydoun MA, Gamaldo AA, Canas JA, Beydoun HA, Shah MT, McNeely JM, et al. Serum Nutritional Biomarkers and Their Associations with Sleep among US Adults in Recent National Surveys. *PLOS ONE*. 2014; 9(8):e103490. <https://doi.org/10.1371/journal.pone.0103490> PMID: 25137304

55. Huang W, Shah S, Long Q, Crankshaw AK, Tangpricha V. Improvement of Pain, Sleep, and Quality of Life in Chronic Pain Patients With Vitamin D Supplementation. *The Clinical Journal of Pain*. 2013; 29(4):341–7. <https://doi.org/10.1097/AJP.0b013e318255655d> PMID: 22699141
56. Kim JH, Chang JH, Kim DY, Kang JW. Association Between Self-Reported Sleep Duration and Serum Vitamin D Level in Elderly Korean Adults. *Journal of the American Geriatrics Society*. 2014; 62(12):2327–32. <https://doi.org/10.1111/jgs.13148> PMID: 25516029
57. Mason C, de Dieu Tapsoba J, Duggan C, Wang C-Y, Korde L, McTiernan A. Repletion of vitamin D associated with deterioration of sleep quality among postmenopausal women. *Preventive Medicine*. 2016; 93:166–70. <https://doi.org/10.1016/j.ypmed.2016.09.035> PMID: 27687537
58. Massa J, Stone KL, Wei EK, Harrison SL, Barrett-Connor E, Lane NE, et al. Vitamin D and Actigraphic Sleep Outcomes in Older Community-Dwelling Men: The MrOS Sleep Study. *Sleep*. 2015; 38(2):251–7. <https://doi.org/10.5665/sleep.4408> PMID: 25581929
59. Song BM, Kim HC, Rhee Y, Youm Y, Kim CO. Association between serum 25-hydroxyvitamin D concentrations and depressive symptoms in an older Korean population: A cross-sectional study. *Journal of Affective Disorders*. 2016; 189:357–64. <https://doi.org/10.1016/j.jad.2015.09.043> PMID: 26476420
60. Darling AL, Skene DJ, Lanham-New SA. Preliminary evidence of an association between vitamin D status and self-assessed sleep duration but not overall sleep quality: results from the D-FINES study of South Asian and Caucasian pre- and post-menopausal women living in Southern England. *Proceedings of the Nutrition Society*. 2011; 70(OCE3):E88. Epub 08/15. <https://doi.org/10.1017/S0029665111001285>.
61. Darling A, Hart K, Middleton B, Berry J, Lanham-New S, Skene D, editors. Lack of association between serum 25-hydroxyvitamin D and sleep quality. *The 21st Vitamin D Workshop*; 2018; Barcelona, Spain 2018.
62. Doo M. The Association between Sleep Duration and 25-Hydroxyvitamin D Concentration with Obesity in an Elderly Korean Population: A Cross-Sectional Study. *Nutrients*. 2018; 10(5):575. <https://doi.org/10.3390/nu10050575> PMID: 29738471
63. Grandner MA, Jackson N, Gerstner JR, Knutson KL. Dietary nutrients associated with short and long sleep duration. Data from a nationally representative sample. *Appetite*. 2013; 64:71–80. <https://doi.org/10.1016/j.appet.2013.01.004> PMID: 23339991
64. Piovezan RD, Hirotsu C, Feres MC, Cintra FD, Andersen ML, Tufik S, et al. Obstructive sleep apnea and objective short sleep duration are independently associated with the risk of serum vitamin D deficiency. *PLOS ONE*. 2017; 12(7):e0180901. <https://doi.org/10.1371/journal.pone.0180901> PMID: 28686746
65. Gong QH, Li SX, Li H, Chen Q, Li XY, Xu GZ. 25-Hydroxyvitamin D Status and Its Association with Sleep Duration in Chinese Schoolchildren. *Nutrients*. 2018; 10(8):1013. <https://doi.org/10.3390/nu10081013>.
66. Gao Q, Kou T, Zhuang B, Ren Y, Dong X, Wang Q. The Association between Vitamin D Deficiency and Sleep Disorders: A Systematic Review and Meta-Analysis. *Nutrients*. 2018; 10(10):1395. <https://doi.org/10.3390/nu10101395> PMID: 30275418.
67. Musiol IM, Stumpf WE, Bidmon HJ, Heiss C, Mayerhofer A, Bartke A. Vitamin D nuclear binding to neurons of the septal, striatal and amygdaloid area in the siberian hamster (*Phodopus sungorus*) brain. *Neuroscience*. 1992; 48(4):841–8. [https://doi.org/10.1016/0306-4522\(92\)90272-4](https://doi.org/10.1016/0306-4522(92)90272-4) PMID: 1321365
68. Gominak SC, Stumpf WE. The world epidemic of sleep disorders is linked to vitamin D deficiency. *Medical Hypotheses*. 2012; 79(2):132–5. <https://doi.org/10.1016/j.mehy.2012.03.031> PMID: 22583560
69. Khan H, Kunutsor S, Franco OH, Chowdhury R. Vitamin D, type 2 diabetes and other metabolic outcomes: a systematic review and meta-analysis of prospective studies. *Proceedings of the Nutrition Society*. 2012; 72(1):89–97. Epub 10/31. <https://doi.org/10.1017/S0029665112002765> PMID: 23107484
70. Mason RS, Rybchyn MS, Abboud M, Brennan-Speranza TC, Fraser DR. The Role of Skeletal Muscle in Maintaining Vitamin D Status in Winter. *Current Developments in Nutrition*. 2019. <https://doi.org/10.1093/cdn/nzz087> PMID: 31598576
71. McCarter SJ, St Louis EK, Sandness DJ, Duwell EJ, Timm PC, Boeve BF, et al. Diagnostic REM sleep muscle activity thresholds in patients with idiopathic REM sleep behavior disorder with and without obstructive sleep apnea. *Sleep Med*. 2017; 33:23–9. Epub 2017/04/30. <https://doi.org/10.1016/j.sleep.2016.03.013> PMID: 28449901; PubMed Central PMCID: PMC5412719.
72. Dattilo M, Antunes HKM, Nunes-Galbes NM, Monico-Neto M, Souza HS, Quaresma M, et al. Effects of Sleep Deprivation on the Acute Skeletal Muscle Recovery after Exercise. *Med Sci Sports Exerc*. 2019. Epub 2019/08/31. <https://doi.org/10.1249/mss.0000000000002137> PMID: 31469710.

73. Mah CD, Mah KE, Kezirian EJ, Dement WC. The effects of sleep extension on the athletic performance of collegiate basketball players. *Sleep*. 2011; 34(7):943–50. Epub 2011/07/07. <https://doi.org/10.5665/SLEEP.1132> PMID: 21731144; PubMed Central PMCID: PMC3119836.
74. Monico-Neto M, Dattilo M, Ribeiro DA, Lee KS, de Mello MT, Tufik S, et al. REM sleep deprivation impairs muscle regeneration in rats. *Growth factors (Chur, Switzerland)*. 2017; 35(1):12–8. Epub 2017/04/19. <https://doi.org/10.1080/08977194.2017.1314277> PMID: 28415893.
75. Yang J, Qiu H, Li H, Zhang Y, Tao X, Fan Y. Body Mass Index, Waist Circumference and Cut-Off Points for Metabolic Syndrome in Urban Residents in Ningxia. *Open Journal of Endocrine and Metabolic Diseases*. 2015; 5(12):163.
76. Liu Y, Tong G, Tong W, Lu L, Qin X. Can body mass index, waist circumference, waist-hip ratio and waist-height ratio predict the presence of multiple metabolic risk factors in Chinese subjects? *BMC Public Health*. 2011; 11(1):35. <https://doi.org/10.1186/1471-2458-11-35> PMID: 21226967
77. Wang F, Wu S, Song Y, Tang X, Marshall R, Liang M, et al. Waist circumference, body mass index and waist to hip ratio for prediction of the metabolic syndrome in Chinese. *Nutrition, Metabolism and Cardiovascular Diseases*. 2009; 19(8):542–7. <https://doi.org/10.1016/j.numecd.2008.11.006> PMID: 19188050
78. Guan X, Sun G, Zheng L, Hu W, Li W, Sun Y. Associations between metabolic risk factors and body mass index, waist circumference, waist-to-height ratio and waist-to-hip ratio in a Chinese rural population. *Journal of Diabetes Investigation*. 2016; 7(4):601–6. <https://doi.org/10.1111/jdi.12442> PMID: 27181937
79. Barceló A, Esquinas C, Piérola J, De la Peña M, Sánchez-de-la-Torre M, Montserrat JM, et al. Vitamin D Status and Parathyroid Hormone Levels in Patients with Obstructive Sleep Apnea. *Respiration*. 2013; 86(4):295–301. <https://doi.org/10.1159/000342748> PMID: 23154407
80. Erden ES, Genc S, Motor S, Ustun I, Ulutas KT, Bilgic HK, et al. Investigation of serum bisphenol A, vitamin D, and parathyroid hormone levels in patients with obstructive sleep apnea syndrome. *Endocrine*. 2014; 45(2):311–8. <https://doi.org/10.1007/s12020-013-0022-z> PMID: 23904340
81. Goswami U, Ensrud KE, Paudel ML, Redline S, Schernhammer ES, Shikany JM, et al. Vitamin D Concentrations and Obstructive Sleep Apnea in a Multicenter Cohort of Older Males. *Annals of the American Thoracic Society*. 2016; 13(5):712–8. <https://doi.org/10.1513/AnnalsATS.201507-440OC> PMID: 26845389.
82. Kerley CP, Hutchinson K, Bolger K, McGowan A, Faul J, Cormican L. Serum Vitamin D Is Significantly Inversely Associated with Disease Severity in Caucasian Adults with Obstructive Sleep Apnea Syndrome. *Sleep*. 2016; 39(2):293–300. <https://doi.org/10.5665/sleep.5430> PMID: 26414899
83. Kerley CP, Hutchinson K, Bramham J, McGowan A, Faul J, Cormican L. Vitamin D Improves Selected Metabolic Parameters but Not Neuropsychological or Quality of Life Indices in OSA: A Pilot Study. *Journal of Clinical Sleep Medicine: JCSM: Official Publication of the American Academy of Sleep Medicine*. 2017; 13(1):19–26. <https://doi.org/10.5664/jcsm.6378> PMC5181609. PMID: 27707440
84. Liguori C, Izzi F, Mercuri NB, Romigi A, Cordella A, Tarantino U, et al. Vitamin D status of male OSAS patients improved after long-term CPAP treatment mainly in obese subjects. *Sleep Medicine*. 2017; 29:81–5. <https://doi.org/10.1016/j.sleep.2016.08.022> PMID: 27964863
85. Liguori C, Romigi A, Izzi F, Mercuri NB, Cordella A, Tarquini E, et al. Continuous Positive Airway Pressure Treatment Increases Serum Vitamin D Levels in Male Patients with Obstructive Sleep Apnea. *Journal of Clinical Sleep Medicine: JCSM: Official Publication of the American Academy of Sleep Medicine*. 2015; 11(6):603–7. <https://doi.org/10.5664/jcsm.4766> PMC4442220. PMID: 25766695
86. Mete T, Yalcin Y, Berker D, Ciftci B, Guven SF, Topaloglu O, et al. Obstructive sleep apnea syndrome and its association with vitamin D deficiency. *Journal of Endocrinological Investigation*. 2013; 36(9):681–5. <https://doi.org/10.3275/8923> PMID: 23558409
87. Terzi R, Yilmaz Z. Bone mineral density and changes in bone metabolism in patients with obstructive sleep apnea syndrome. *Journal of Bone and Mineral Metabolism*. 2016; 34(4):475–81. <https://doi.org/10.1007/s00774-015-0691-1> PMID: 26204846
88. Autier P, Boniol M, Pizot C, Mullie P. Vitamin D status and ill health: a systematic review. *The lancet Diabetes & endocrinology*. 2014; 2(1):76–89. [https://doi.org/10.1016/S2213-8587\(13\)70165-7](https://doi.org/10.1016/S2213-8587(13)70165-7).
89. Anglin RES, Samaan Z, Walter SD, McDonald SD. Vitamin D deficiency and depression in adults: systematic review and meta-analysis. *British Journal of Psychiatry*. 2013; 202(2):100–7. Epub 01/02. <https://doi.org/10.1192/bjp.bp.111.106666>.
90. Irwin MR, Olmstead R, Carroll JE. Sleep Disturbance, Sleep Duration, and Inflammation: A Systematic Review and Meta-Analysis of Cohort Studies and Experimental Sleep Deprivation. *Biological Psychiatry*. 2016; 80(1):40–52. <https://doi.org/10.1016/j.biopsych.2015.05.014> PMID: 26140821