

8-1-2020

## The relationship between sleep duration and mood in adolescents: A systematic review and meta-analysis

Michelle A. Short  
*Flinders University*

Stephen A. Booth  
*CQUniversity Australia*

Omar Omar  
*University of Birmingham*

Linda Ostlundh  
*United Arab Emirates University*

Teresa Arora  
*Zayed University, [teresa.arora@zu.ac.ae](mailto:teresa.arora@zu.ac.ae)*

Follow this and additional works at: <https://zuscholars.zu.ac.ae/works>



Part of the [Medicine and Health Sciences Commons](#)

---

### Recommended Citation

Short, Michelle A.; Booth, Stephen A.; Omar, Omar; Ostlundh, Linda; and Arora, Teresa, "The relationship between sleep duration and mood in adolescents: A systematic review and meta-analysis" (2020). *All Works*. 3561.

<https://zuscholars.zu.ac.ae/works/3561>

This Article is brought to you for free and open access by ZU Scholars. It has been accepted for inclusion in All Works by an authorized administrator of ZU Scholars. For more information, please contact [scholars@zu.ac.ae](mailto:scholars@zu.ac.ae).

# The relationship between sleep duration and mood in adolescents: A systematic review and meta-analysis

Short, Michelle A.; Booth, Stephen A.; Omar, Omar; Ostlundh, Linda; Arora, Teresa

DOI:

[10.1016/j.smr.2020.101311](https://doi.org/10.1016/j.smr.2020.101311)

License:

Creative Commons: Attribution-NonCommercial-NoDerivs (CC BY-NC-ND)

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Short, MA, Booth, SA, Omar, O, Ostlundh, L & Arora, T 2020, 'The relationship between sleep duration and mood in adolescents: A systematic review and meta-analysis', *Sleep Medicine Reviews*, vol. 52, 101311. <https://doi.org/10.1016/j.smr.2020.101311>

[Link to publication on Research at Birmingham portal](#)

## General rights

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

## Take down policy

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# Journal Pre-proof

The Relationship Between Sleep Duration and Mood in Adolescents: A systematic review and meta-analysis

Michelle A. Short, Stephen A. Booth, Omar Omar, Linda Ostlundh, Teresa Arora



PII: S1087-0792(20)30054-X

DOI: <https://doi.org/10.1016/j.smr.2020.101311>

Reference: YSMRV 101311

To appear in: *Sleep Medicine Reviews*

Received Date: 14 November 2019

Revised Date: 30 January 2020

Accepted Date: 31 January 2020

Please cite this article as: Short MA, Booth SA, Omar O, Ostlundh L, Arora T, The Relationship Between Sleep Duration and Mood in Adolescents: A systematic review and meta-analysis, *Sleep Medicine Reviews*, <https://doi.org/10.1016/j.smr.2020.101311>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Elsevier Ltd. All rights reserved.

# The Relationship Between Sleep Duration and Mood in Adolescents: A systematic review and meta-analysis

Running head: Sleep Duration and Mood in Adolescents

Michelle A. Short<sup>1</sup>, Stephen A. Booth<sup>2</sup>, Omar Omar<sup>3</sup>, Linda Ostlundh<sup>4</sup> & Teresa Arora<sup>5</sup>

1. Flinders University, Adelaide, South Australia, AUSTRALIA.
2. Central Queensland University, Rockhampton, Queensland, AUSTRALIA.
3. University of Birmingham, Birmingham, UNITED KINGDOM.
4. United Arab Emirates University, Al Ain, UNITED ARAB EMIRATES.
5. Zayed University, Abu Dhabi, UNITED ARAB EMIRATES.

The authors have no conflicts of interest to declare.

Correspondence to:

Dr Teresa Arora  
Assistant Professor  
Zayed University  
College of Natural and Health Sciences  
Department of Psychology  
Office FF1-1-12  
P O Box 144534  
Abu Dhabi  
United Arab Emirates  
Tel: +971 544 497 305  
Email: Teresa.Arora@zu.ac.ae

## SUMMARY

Insufficient sleep has been argued to result in deleterious changes to mood in adolescents and offers promise as a modifiable risk factor. A systematic review of the literature regarding sleep duration and mood in adolescents was conducted using the academic databases PsycINFO, PubMed, Medline, Scopus, and EMBASE to identify relevant literature. Seventy-four studies, including 361,505 adolescents were sourced out of the 1534 references identified, 73 of which were appropriate for meta-analysis. Pooled results indicated that less sleep was associated with a 55% increase in the likelihood of mood deficits. Positive mood showed the largest relationship with sleep duration, followed by anger, depression, negative affect and anxiety. Effect sizes also varied according to study design, how sleep was operationalised, and geographical region, but not according to the inclusion of covariates. Sleep duration has a significant negative impact on a range of mood states in healthy adolescents. These effects were witnessed across all geographical regions, highlighting that sleep is a universal and modifiable risk factor for preventing mood deficits in this at-risk population.

**Keywords:** adolescence, teens, sleep duration, mood, affect, depression, anxiety.

## INTRODUCTION

Despite being identified as a strong correlate of adolescent mental and physical health, academic performance, and relationship quality (1-3), literature examining the causal contributors to adolescent mood is limited which, in turn, hinders the development of intervention strategies. Understanding causal contributors to mood is paramount for adolescents, who run a heightened risk of developing a mood disorder as the result of the psychological, social and physiological shifts that occur at this developmental stage (4). One survey of over 10,000 U.S. adolescents aged 13 to 18 years found a lifetime prevalence of anxiety disorders of 31.9% and a lifetime prevalence of mood disorders of 14.3% (5). Adolescence is a vulnerable period during which the onset of many mood disorders occur (6, 7). Mood disorders such as depression and anxiety often have a chronic and recurrent course, with earlier age of onset associated with poorer educational, social and quality of life outcomes (4). Indeed, mental illness conveys one of the largest disease burdens of all health conditions in terms of both mortality and morbidity (8), with depression projected to be the second leading cause of disease burden worldwide by 2020 (9).

One factor posited to have a causal relationship with mood is sleep (10-12). Research has identified sufficient sleep as a contributor to optimal mood and the ability to better regulate emotions (12-14). Despite the extant evidence highlighting the importance of sleep across a range of psychological outcomes, the overwhelming majority of adolescents obtain insufficient sleep (15-17). For example, a National Sleep Foundation Poll reported that more than 87% of U.S. high school students obtain less than the recommended hours of sleep (18). It is recommended that adolescents obtain between 8 and 10 hours of sleep per night (14, 19, 20), however, bedtimes delay across adolescence while the need to rise for school remains the same or moves even earlier, thus limiting sleep (21, 22). Several factors unique to

adolescence contribute to short sleep, including the puberty-related delay in circadian timing, slower accumulation of homeostatic sleep pressure in the evening, earlier school start times, diminished parental involvement in setting bedtimes, and increased autonomy over the use of electronic devices

Considering the prevalence of shortened sleep in adolescents, and the importance of mood as a contributor to mental health, it is imperative to consider the effects of insufficient sleep on adolescent mood. A recent systematic review and meta-analysis examined the relationship between sleep and depression in adolescents across 23 studies (23). The meta-analysis revealed that depressed adolescents experienced more wakefulness in bed through taking longer to fall asleep and more wakefulness during the night, and they reported more subjective sleep disturbances. Examination of the longitudinal trajectories of both sleep and depression suggested that sleep disturbance was a precursor for the development of depression (23). The present review expands on this work by examining mood more broadly, to include a range of both positive and negative mood states, rather than focusing solely on mental health diagnoses. While positive and negative mood are sometimes conceptualised as occurring at either end of the same spectrum of mood, they can also be regarded as independent constructs. As such, including discrete mood states acknowledges that, (a) mood occurs on a spectrum broader and more nuanced than simply the presence or absence of mental illness, and (b) that positive moods also play an important role in mental health.

In addition, while the relationship between sleep and mood is relatively well established in clinical samples, less is known about whether mood changes are induced in otherwise healthy adolescents when subjected to shorter sleep durations. As such, the present review includes only non-clinical samples of adolescents. Establishing a causal relationship between sleep and mood in healthy adolescents will support the development of mainstream interventions regarding sleep that may be implemented by schools, policymakers and parents

to assist in the healthy development of adolescents and foster an early intervention approach to sleep and mental health. The present review examines the academic literature to understand the relationship between sleep duration and mood in adolescents. We also investigate whether the relationship between sleep duration and mood varies according to factors such how sleep is measured, which mood state is assessed, geographical location, study design, and the inclusion of covariates.

## METHOD

A literature search was used to identify original studies examining the relationship between sleep duration and mood in adolescents. Relevant literature was sourced using the academic databases PsycINFO, PubMed, Medline, Scopus, and EMBASE in December 2018. The following basic search string was developed through pre-searches in PubMed and applied in all four databases: ((happiness OR suicidal OR antagonism OR apathy OR oppositionality OR euphori\* OR depress\* OR tense\* OR tension OR irrita\* OR annoyed OR hostile\* OR hate\* OR frustrat\* OR rage OR moods OR mood OR moody OR anxious\* OR temper OR emotion\* OR anxiety OR distress\* OR anger\* OR angry OR happy\* OR sad OR sadness OR confus\* OR upset) AND ("sleep duration" OR "sleep deprivation" OR "sleep loss" OR "total sleep" OR "sleep restriction") AND (adolescen\* OR youth\* OR teenager\* OR teen OR teens)). Searches occurred on December 14 and 17, 2018. All terms were searched with a combination of the fields "Title", "Abstract" and "MeSH/Thesaurus" (when available) for best possible search precision. No filters or limitations except for "English language only" were applied to ensure inclusion of pre-indexed materials. A total of 2,162 references were identified through the search and uploaded to the systematic review software Covidence for de-duplication, blinded screening and extraction. A detailed search log



including all search strings, results and notes is available in Appendix A. This review is registered on Prospero, ID CRD42017068617.

Studies were included if they were primary peer-reviewed journal articles or articles in press, included participants aged 10 to 19 years, as per the World Health Organisation definition of adolescence (24), examined the relationship between sleep duration and mood, and were published in English. Studies that did not report the relationship between sleep and mood in a healthy population (i.e., included clinical population only), or that included pharmacological interventions, were excluded. No exclusion criteria were applied regarding the date of publication to capture all relevant published original research. Data extracted from each study included authors, year of publication, sample size, region (Asia, Australia and New Zealand, Europe, or North America), age range, proportion of male participants, how sleep duration and mood were operationalised, study design, and effect size.

Forty-two, out of the 74 included studies, utilised a cross-sectional design. As such, study quality was assessed for selection bias, information bias, and confounding, which are the three primary sources of bias in these studies (16, 25). Selection bias was assessed using three criteria: Were inclusion criteria given and applied uniformly? Did the sampling strategy achieve a sample representative of the target population? Was the response rate  $\geq 80\%$ ? Information bias refers to the reliability and validity of study measures. As subjective measures contain a higher risk of bias due to factors such as social desirability responding and inaccuracy, information bias was assessed using one criterion: Was the IV (sleep duration) measured objectively? Finally, the risk of confounding variables was assessed using one criterion: Did the study assess and statistically adjust for confounding variables?

### *Statistical Analyses*

The effect sizes were coded to represent the odds of mood deficits with shorter sleep durations; thus, higher values represent worse mood outcomes (for example, greater anger, anxiety, depressed mood and negative affect and/or decreased positive mood affect) with less sleep. Because the selected studies were carried out in different settings and evidence of high level of heterogeneity, random-effects model was employed to calculate the overall effect from effect sizes. Between-study heterogeneity was assessed using I-square ( $I^2$ ) test (26). A priori subgroup analyses were performed to explore the impact on the effect size. Five subgroup analyses were performed to investigate if the relation between sleep duration and mood was different: 1) Region - Asia, Australia/New Zealand, Europe and North America, 2) Study design – Experimental, longitudinal or cross-sectional, 3) Covariate adjustment – demographics (such as sex or age), others (such as family income or parental years of education, snoring), none, 4) Mood – anger, depressed mood, anxiety, positive effect and negative effect, 5) Sleep operationalisation – actigraphy, sleep diary, questionnaires or PSG. While anger, depressed mood and anxiety could all be classified as aspects of negative affect, because it was possible to include these more specific classifications of negative affect, we chose to do this where possible to allow for a more fine-grained evaluation across different mood states.

Heterogeneity between subgroups was evaluated using random-effect model. Two-sided p-values  $\leq 0.05$  were considered statistically significant for all tests. To calculate the overall effect size, same studies reporting several mood effects were combined by intrastudy meta-analyses to create independent effect sizes required for the meta-analysis (27).

Publication bias was assessed visually using a funnel plot.

## RESULTS

Results of the literature search revealed 74 studies including 361,505 adolescents. The PRISMA flow chart detailing the systematic screening and identification of studies is shown in Figure 1. Studies included mood states such as positive affect (N=10), negative affect (N=11), depressed mood (N=50), anger (N=5), anxiety (N=26), confusion (N=3), irritability (N=3), oppositionality (N=1), stress (N=1), mood (N=1), happiness (N=2), sadness (N=1) and nervousness (N=2). Of the 74 studies, 44 used subjective self-report to measure sleep duration, nine used sleep diaries, 16 used actigraphy, and five used polysomnography (PSG). Examination of study design reveals varied approaches, with 42 cross-sectional, 19 longitudinal and 13 experimental studies identified. The majority of included studies (N=29) were conducted in North America, with 17 studies from Europe, 17 from Asia and 11 from Australia. A summary of the included studies is provided in Table 1.

Insert Figures 1 and 2 here

Meta-analytic results are shown in Figure 2. Overall, results indicated that shorter sleep durations were associated with a 55% increased risk of mood deficits,  $OR = 1.55$ ,  $p < .001$ . Effects sizes were significantly heterogeneous between studies,  $I^2 = 97.9$ ,  $p < .001$ . To examine whether the effect of sleep duration on mood varied depending on the type of mood state assessed, study design, geographical region, inclusion of covariates, or how sleep was operationalised, further meta-analyses were conducted. Effect sizes for each subgroup are provided in Table 2.

Effect sizes significantly varied between mood states,  $Q(4) = 88.23$ ,  $p < .001$ . Mood states that were examined in fewer than 4 studies were not included as effect size estimates are less reliable when fewer than 4 studies are included. Positive affect showed the largest deficits following shorter sleep, followed by anger, depression, negative affect and anxiety. There was significant heterogeneity among effect sizes according to how sleep was operationalised,  $Q(3) = 57.01$ ,  $p < .001$ . Studies measuring sleep using polysomnography

had the largest effect sizes on mood, followed by questionnaires, actigraphy and sleep diaries. Effect sizes varied according to study design,  $Q(2) = 144.28, p < .001$ . Experimental and longitudinal studies revealed larger effect sizes between sleep duration and mood than cross-sectional studies. There was significant heterogeneity among effect sizes according to geographical region,  $Q(3) = 46.98, p < .001$ , with North American studies reporting larger effect sizes, followed by European studies, Australian and New Zealand studies and Asian studies. Effect sizes were not significantly heterogeneous according to the covariates included,  $Q(2) = 5.58, p = .06$ . Results of the risk of bias assessment is shown in Figure 3. A funnel plot of effect sizes is shown in Figure 4 and appears symmetrical, indicating no significant publication bias.

Insert Figure 3 and 4 here

Table 1. Summary of articles identified in the systematic review. Note: S = subjective, O = objective, P = prospective, C = cross-sectional, L = longitudinal, E = experimental, TST = total sleep time, TIB = time in bed.

First Author (Year)	N	Country	Slee p	Mood variable	Study design	Summary
Barnes & Meldrum (2014) (28)	563	US	S	Depressed mood	C	Significant negative association between sleep duration and depressed mood
Bauducco et al., (2016) (29)	1,057	Sweden	S	Depressed mood, anger, anxiety	C	Significant dose-dependent effect of sleep duration on depressed mood, anger and anxiety among 12-13yo and 14-16yo
Baum et al., (2014) (30)	50	US	O	Anxiety, depression, anger, fatigue, confusion, vigour	E	Adolescents reported more anxiety, anger, fatigue, confusion and less vigour with sleep restricted to 6.5h TIB for 5 nights compared to 10h TIB for 5 nights. No changes to depression were found
Beebe et al., (2008) (31)	19	US	O	Oppositionality/irritability	E	Parents, but not adolescents, reported that adolescents were more irritable/oppositional during short vs extended sleep
Bei et al., (2013) (32)	9	Australia	O	Generalized anxiety,	L	The small increase in TST post-intervention was associated with a small increase in general anxiety
Bei et al., (2017) (33)	146	Australia	O	Negative mood (depressed mood and anxiety)	C	No significant association between negative mood (depressed mood and anxiety) and TIB during school holidays
Boergers et al., (2014) (34)	197	US	S	Depressed mood	L	Depressed mood decreased after a 30-minute delay to school start times
Bonnar et al., (2015) (35)	141	Australia	S	Depressed mood	L	Depressed mood decreased following a sleep education program after an increase in school night TST of 27 minutes
Chan et al., (2018) (36)	82	Hong Kong	S	Depression, anxiety, stress	L	Depression, anxiety and stress decreased following a delay in school start time by one hour and increase in TST.
Chue et al., (2018) (37)	89	US	O	Positive affect, negative affect	L	Higher objectively measured TST was associated with better mood recovery following stress for positive and negative affect
Conklin et al., (2018) (38)	3,017	Canada	S	Depressed mood	L	Chronic sleep deprivation was significantly correlated with depression in females but not males
Dewald-Kaufmann et al., (2014) (39)	55	Netherlands	O	Depressed Mood	E	Adolescents in the sleep extension group obtained more sleep and their depressive symptoms diminished significantly compared to controls
Diaz-Morales (2016) (40)	1406	Spain	S	Anxiety	C	There was a significant negative relationship between TIB and anxiety on school nights but not weekends
Doane & Thurston (2014) (41)	78	US	O	Positive affect, negative affect	C	No significant associations were found between TST and positive or negative affect
Doane et al., (2015) (42)	82	US	O	Anxiety, depressed	L	No significant association between sleep duration at Time 1 and depressed

				mood		mood or anxiety one year later
El-Sheikh et al., (2016) (43)	252	US	O	Depressed mood, anxiety	C	No significant associations between sleep duration and depressed mood or anxiety
Fan et al., (2017) (44)	1,573	China	S	Depressed mood (Dichotomised Y/N)	L	After controlling for depression and covariates at Time 1, sleep duration at T1 did not significantly predict depression at T2
Fredrikson et al., (2004) (45)	2,259	US	S	Depressed mood	L	Students who slept less over time experienced more depressed mood
Fuligni et al., (2017) (14)	419	US	P	Depressed mood and anxiety	L	A non-linear association existed between TST and psychological distress the following day, with both too much and too little sleep were associated with next-day distress.
Hall Brown (2008) (46)	35	U.S.	P	General mood	E	Sleep duration, but not general mood, was better on weekdays than weekends
Hasler (2008) (47)	56	U.S.	NS	Anxiety	E	Sleep extended by one hour per night for 3 consecutive nights did not result in decreased anxiety relative to non-extended group
Hyakutake et al., (2016) (48)	409	Japan	S	Depressed mood	C	U-shaped relationship between sleep duration and depressed mood
Itani et al., (2016) (49)	94,777	Japan	S	Anger	C	Adolescents who slept less than 5h per night were significantly more likely to report anger
Josev et al., (2017) (50)	136	Australia	O	Anxiety	C	TST was not significantly associated with anxiety in healthy controls
Kalak et al., (2012) (51)	80	Switzerland	O	Depressed mood	C	No significant association between objective TST and depressed mood
Kang et al., (2014) (52)	4,145	Korea	S	Depressed Mood	C	Participants who slept less than 7h on school nights reported significantly higher levels of depressed mood
Kelly & El-Sheikh, (2014) (53)	113	U.S.	O	Depressed Mood, anxiety	L	Sleep minutes at T2 (10 years) significantly predicted depressed mood at T3 (13 years) but not anxiety at T3
Kuo et al., (2015) (54)	246	U.S.	P	Depressed mood	C*	No relationship between sleep duration and depressed mood
Lazaratou et al., (2013) (55)	696	Greece	S	Anxiety (state and trait)	C	Sleep duration had a small but significant association with state and trait anxiety
Lee et al., (2016) (56)	3,785	Korea	S	Depressed mood	C	Sleep duration on both weekdays (M-F) and weekends (Sa-Su) had small but significant negative associations with depressed mood
Lee et al., (2012) (57)	8,010	Korea	S	Depressed mood	C	Sleep duration on both weekdays and weekends were significantly and negatively associated with depressed mood
Lee (2017) (58)	31,407	Korea	S	Depressed mood	C	TST was negatively correlated with feelings of sadness and hopelessness
Lehto et al., (2014) (59)	439	Finland	S	Depressed mood	C	Sleep duration on weekdays and weekends showed a small but significant negative association with depressed mood
Lemola et al., (2011) (60)	190	Germany Austria Switzerland	S	Depressed mood	C	No significant association between sleep duration and mood on weekdays not weekends

Lemola et al., (2015) (61)	362	Switzerland	S	Depressed mood	C	Sleep duration was negatively associated with depressed mood
Lin et al., (201) (62)	2,472	Taiwan	S	Emotional wellbeing (Happiness & Depressed mood)	L	Adolescents who slept less than 6h per night were significantly more likely to have low levels of emotional well-being
Liu & Zhou (2002) (63)	1,359	China	S	Anxious/depressed mood subscale	C	Adolescents sleeping less than 8h had significantly higher odds of depressed mood and anxiety than those with 9h or more.
Lo et al., (2017) (64)	57	Singapore	O	Positive affect, negative affect	E	Sleep restriction lead to reduced positive affect but no change to negative affect
Lo et al., (2016) (65)	56	Singapore	O	Positive affect, negative affect	E	Sleep restriction diminished positive affect but not negative affect
Lovato et al., (2017) (66)	138	Australia	S	Depressed mood	L	Sleep duration at T1 did not predict depressed mood 1 year later, however, T1 TST was associated with depressed mood at T1
Maume (2017) (67)	974	U.S.	S	Depressed Mood	L	Sleep duration had a negative correlation with depressive symptoms.
Mazzer et al., (2018) (68)	951	Sweden	S	Depressed mood, anxiety	L	Sleep duration had a negative correlation with psychological distress
McGlinchey et al., (2011) (69)	38	U.S.	O	Computerised and human rated positive and negative emotions	E	Compared to when well-rested, sleep restricted to 2h decreased positive emotional expression but did not affect negative emotion using computer text analysis, and less positive emotional expression and more negative emotion using human raters
McMakin et al., (2016) (70)	64	US	O	Positive affect, negative affect	E	Experimental sleep restriction to 4h TIB led to decreased positive affect and increased negative affect compared to affect following a 10h sleep opportunity
Moore et al., (2009) (71)	247	US	O	Anxiety, depressed mood	C	Sleep duration was not significantly associated with depressed mood or anxiety
Mullin et al., (2017) (72)	17	US	P	Morning Anxiety	L	No significant relationship between sleep duration and anxiety morning
Norell-Clarke & Hagquist (2018) (73)	15,221	Sweden	S	Sadness, nervousness, irritation	C	Individuals sleeping less than 8 hours were more likely to report feeling sad, nervous and irritated than those sleeping more than 8 hours
Nowakowski et al., (2016) (74)	1,042	US	S	Depressed mood	L	After controlling for depressed mood at T1, sleep duration at T2 significantly predicted depressed mood one year later
Nuutinen et al., (2014) (75)	4,985	Denmark, Finland, France	S	Negative affect (feeling low, irritable, nervous)	C	Significant negative associations were found between sleep duration and negative affect among all adolescents except for Danish males.
Oginska & Pokorski, (2006) (76)	191	Poland	S	Negative mood	C	No association between TIB on school nights and negative mood
Ojio t al., (2016) (13)	15,637	Japan	S	Depression and	C	Depression/anxiety symptoms were the lowest in males and females who

				anxiety (above/below cut-off)		slept 8.5–9.5 h and 7.5–8.5 h, respectively and worse for both shorter and longer sleepers
Owens et al., (2010) (77)	201	US	S	% reporting being at least somewhat unhappy or depressed, % reporting feeling irritated or annoyed, Depressed mood	E	A 30m delay to school start time led to a 45m increase in TST on school nights, less depressed mood, and a significantly lower proportion of adolescents who reported feeling at least somewhat unhappy or depressed or feeling irritated or annoyed
Pallesen et al., (2010) (78)	1,279	Norway	S	Anxiety, Depression	C	Short sleepers (<7h) had significantly higher depressed mood but not anxiety when compared to normal sleepers (≥7h)
Pasch et al., (2010) (79)	242	US	S	Depressed mood	C	Depressed mood was negatively associated with depressed mood and school night sleep duration, but not weekend sleep duration
Randler & Weber, (2015) (80)	219	Germany	S	Positive affect, Negative affect	C	Negative, but not positive, affect was significantly associated with TST the prior night. Neither positive or negative affect were related to habitual sleep duration
Raniti et al., (2017) (81)	695	Australia	S	Depressed mood	C	School night sleep duration was negatively associated with depressed mood
Reddy et al., (2017) (82)	42	US	O	Positive affect Negative affect State Anxiety Trait Anxiety	E	Adolescents whose sleep was restricted to 4h TIB for one night reported significantly less positive affect and greater state and trait anxiety, but not negative affect, than adolescents who had 9.5h TIB
Roberts & Duong (2014) (83)	3,134	US	S	Depression symptoms	L	Short sleep was a much larger predictor of major depression than depression symptoms.
Sarchiapone et al., (2014) (84)	11,788	Austria Estonia France Germany Hungary Ireland Israel Italy Romania Slovenia Spain	S	Anxiety	C	School night sleep duration showed a small negative relationship with anxiety
Seo et al., (2017) (85)	26,395	Korea	S	Depressed mood	C	School night sleep duration had a small but significant negative association with depressed mood
Shen et al., (2018) (86)	4,582	Australia	S	Happiness, positive	C	Short sleep was associated with lower happiness and positive affect, and



				affect, negative affect		higher negative affect.
Short & Louca (2015) (87)	12	Australia	O	Depressed mood, anger, confusion, anxiety	E	Depressed mood, anger, confusion and anxiety all significantly increased following one night without sleep
Short et al., (2013a) (88)	385	Australia	P	Depressed mood	C	Significant negative relationship between sleep duration and depressed mood
Short et al., (2013b) (21)	258 <sup>^</sup>	Australia	P	Anxiety	C	Significant negative relationship between sleep duration and anxiety
Singh et al., (2018) (89)	501	India	S	Depressed mood	C	TST was not significantly correlated with depressed mood
Stheneur et al., (2017) (90)	855	France	P	Depressed mood	C	TST was negatively correlated with depressed mood
Suzuki et al., (2011) (91)	99,668	Japan	S	Depression/ anxiety, loss of positive emotion	C	Sleep duration of less than 7h, but not 8h or more, was associated with increased depression/anxiety and loss of positive emotion, while sleep of 8 h or more was associated with greater loss of positive emotion
Van Dyk et al., (2017) (92)	54	US	O	Anxiety, Depressed mood, anger, confusion	E	Extended sleep (average of 72 minutes) resulted in lower anger and confusion.
Wahlstrom et al., (2017) (93)	8,261	US	S	Feeling unhappy, sad, or depressed, and feeling nervous or tense	C	Adolescents who slept longer on school nights were significantly less likely to report feeling unhappy, sad or depressed or nervous or tense
Warner et al., (2008) (94)	307	Australia	S	Depressed mood	C	Significant association between school night TST and depressed mood, with less sleep associated with worse mood
Watson & Brickson, (2018) (95)	65	UK	P	Mood	L	TST negatively correlated with mood
Wolfson & Carskadon, (1998) (22)	2,166	US	S	Depressed mood	C	Adolescents sleeping $\geq 8h15m$ had significantly lower levels of depressed mood than those sleeping $\leq 6h45m$
Yang & Cha (2018) (96)	421	Korea	S	Depressed mood	C	TST negatively correlated with depressed mood
Yip (2015) (97)	146	US	P	Depressed mood, anxiety	C	No significant associations between weekly sleep duration and depressed mood or anxiety

\*Cross-sectional data drawn from a longitudinal study; <sup>^</sup>Subsample of above study

Table 2. Meta-analyses of studies examining the relationship between sleep duration and mood in adolescents. The number of studies for mood states category equal more than (Total N of studies) as some studies included more than one mood state.

	<i>k</i>	<i>OR</i>	<i>Lower limit</i>	<i>Upper limit</i>	<i>p</i>
<b>Overall</b>	<b>73</b>	<b>1.55</b>	<b>1.44</b>	<b>1.67</b>	<b>&lt;.001</b>
<i>Mood State</i>					
Depressed Mood	43	1.62	1.38	1.85	<.001
Anxiety	22	1.41	1.29	1.54	<.001
Anger	9	1.83	1.51	2.15	<.001
Negative Affect	20	1.60	1.49	1.71	<.001
Positive Affect	9	2.02	1.62	2.42	<.001
<i>Sleep Measurement</i>					
Questionnaires	45	1.55	1.44	1.67	<.001
Sleep Diary	9	1.30	1.11	1.49	<.001
Actigraphy	14	1.45	1.28	1.62	<.001
PSG	5	1.70	1.20	2.20	<.001
<i>Region</i>					
Asia	16	1.34	1.19	1.50	<.001
Australia and New Zealand	11	1.42	1.22	1.62	<.001
Europe	17	1.47	1.34	1.60	<.001
North America	29	1.70	1.43	1.98	<.001
<i>Study design</i>					
Experimental	13	1.72	1.48	1.97	<.001
Longitudinal	19	1.71	1.31	2.10	<.001
Cross-sectional	41	1.42	1.31	1.53	<.001
<i>Covariates</i>					
None	45	1.67	1.48	1.85	<.001
Demographics	12	1.44	1.27	1.60	<.001
Others	16	1.28	1.18	1.38	<.001

Note. *k* = no. of studies; OR = Odds Ratio; PSG=polysomnography

## DISCUSSION

The present systematic review and meta-analysis identified 74 original studies, including over 360,000 adolescents, that examined the effect of sleep duration on mood. Of these, 73 studies were suitable for meta-analysis. Shorter sleep duration significantly increased the odds of adolescents experiencing mood deficits by 55%. Mood deficits due to shorter sleep were observed across mood states including depression, anxiety, anger, positive affect and negative affect, indicating that short sleep can lead to a range of mood deficits in otherwise healthy, non-clinical samples of adolescents. While strong research evidence exists to support the casual role of sleep in mood, the important task of estimating the sleep duration for optimal mood is only just emerging (13, 14). Fuligni and colleagues used two weeks of daily estimates of sleep duration and mood to determine the average amount of sleep needed for the lowest levels of next-day depressed mood and anxiety. Estimates revealed that 9-hours sleep per night was needed for optimal mood and that this optimum tended to be greater among younger adolescents and among those with higher levels of psychopathology (14). These estimates concur with estimates of sleep durations for optimal cognitive functioning and with current sleep recommendations (19, 20, 98).

There was significant heterogeneity in effect sizes according to the mood state measured. Shorter sleep durations doubled the odds of adolescents experiencing reduced positive affect, and increased the odds of anger by 83%, depressed mood by 62%, negative affect by 60% and anxiety by 41%. The finding that positive mood is most affected by sleep loss is consistent with recent research (65, 86). For example, Shen and colleagues examined the associations between sleep duration and mood in 4,582 adolescents. They found that the association between sleep loss and mood was stronger for happiness and positive affect than it was for negative affect. Anhedonia, or the reduced ability to experience pleasure from normally pleasurable activities, is a clinically significant symptom and diminishes quality of

life among those experiencing it. These results highlight an important potential mechanism between short sleep and mood disorders through reduced enjoyment or pleasure and elucidates the importance of measuring moods beyond common measures of depression and anxiety (99).

While anhedonia is one mechanism through which sleep loss may affect mood, there are many additional pathways and mechanisms posited to account for this relationship. Sleep loss is known to affect brain regions implicated in mood and emotion regulation. Sleep loss reduces prefrontal activity and reduced functional connectivity between the prefrontal cortex and limbic regions (100). Sleep loss also reduces rapid eye movement (REM) sleep, which is implicated in the processing of emotional memories (101). Finally, sleep loss negatively affects cognitive functioning, many of which processes are needed for affective monitoring, reasoning and emotion regulation (12, 20, 102). For example, the repetitive negative thinking that is a transdiagnostic risk factor for various mental health conditions is argued to be maintained by sleep loss as sleep loss diminishes a person's ability to shift attention away from their repetitive negative thoughts (103).

In addition to differences between mood states, results indicated that effect sizes varied according to the way sleep was operationalised, the design of the study, and geographical region. Studies that utilised an experimental design to manipulate sleep reported the largest effects between sleep duration and mood. This finding is likely to be due, at least in part, to the observation that experimental studies include meaningful manipulations of sleep duration and by having a control condition that included sleep durations within the recommended range. Given the universality of insufficient sleep among adolescents worldwide, cross-sectional and longitudinal studies are likely to include a more restricted range of sleep durations and contain a paucity of adolescents whose sleep falls within the recommended range, thus masking the true effect of sleep duration on mood (104). Effect

sizes from experimental studies are also less likely to be impacted by extraneous variables that are likely confounded with naturalistic sleep loss, such as academic pressure, parental regulation of sleep, and socioeconomic status. Studies using polysomnography to measure sleep duration and mood reported largest effect sizes than those utilising sleep diaries, actigraphy and questionnaires. This may be due to the superior validity of polysomnography and also the improved temporal association between sleep measurement and mood assessment in these studies. Studies using polysomnography typically assessed mood immediately following the objectively measured sleep period, whereas questionnaires ask about “typical” sleep and not the sleep immediately prior to mood assessment. It is important to note, however, that sleep measurement tends to be confounded with study design, as all studies using polysomnography were experimental. Small differences in effect size estimates were observed according to region, with North American studies reporting the largest effect sizes. It is unclear whether these larger effects are due to methodological differences or differences in vulnerability to sleep loss. Cross-cultural studies that directly compare regions using the same methodology are needed to tease apart this finding. Of greater importance was the observation that none of the geographical regions included were spared the negative impacts of shorter sleep durations on mood, with Asia, Europe, North America and Australia and New Zealand studies uniformly similarly finding this relationship. Further studies examining the relationship between sleep and mood in understudied South American, African and Middle eastern countries are needed to understand any potential differences in risk of shorter sleep between countries and/or geographical regions.

The methodological and theoretical implications discussed above provide direction for future research. Firstly, dose-response experimental protocols are required to examine causal relationships between sleep duration and mood. Most of the studies were cross-sectional. Cross-sectional designs are unable to determine the direction of the relationship

between sleep and mood. This is of central importance as research highlights the bi-directional relationship between these variables (13, 99). In addition, cross-sectional studies cannot determine the degree to which confounding factors, such as diet, long-term habitual sleep loss or varying home environments, influence participants' mood. Secondly, an overwhelming emphasis is placed on negative mood states in reviewed studies which is indicative of a research field that has largely focussed on the relationship between sleep loss and the negative mood states of depression and anxiety. The few studies measuring positive mood states, such as positive affect, happiness and vigour derived findings worthy of further investigation as consistent reductions across positive mood states following short sleep were found.

Finally, mood is often conflated with emotion and emotion regulation in reviewed literature and separation of these concepts will be beneficial in future research (12). As all these factors play a key part in the development and perpetuation of clinical disorders such as depression, obsessive-compulsive disorder, anxiety disorders and posttraumatic stress disorder (12), examining explicit links between sleep duration and emotion and emotion regulation, as well as mood, will allow for more accurate prediction of how short sleep may contribute to mood deficits and dysfunction. Similarly, there are multiple aspects of affective functioning that remain underexplored in relation to sleep, including anhedonia, specific types of emotion regulation strategies, and rumination. It is recommended that psychological disorder-specific models be utilised to further examine how sleep loss contributes to the development of mental health disorders among healthy adolescents and how sleep loss maintains these disorders.

## Conclusion

Sleep duration significantly predicted mood deficits on all mood states, including increased depression, anxiety, anger, negative affect and reduced positive affect. This effect was observed across geographical regions, demonstrating that short sleep is truly a universal risk-factor for mood deficits. Fortunately, sleep loss is an inherently *modifiable* risk-factor, providing a target for intervention at individual, family, community and public policy levels. Interventions to increase sleep duration include increasing parental regulation of sleep and technology use, delayed school start times, ensuring academic pressure and out of school hours tutoring does not impede sleep, and maintaining regular sleep and wake times across school nights and weekends. Targeting sleep in this at-risk population will reduce the likelihood of mood deficits transitioning to clinically significant problems and mood disorders.

### **Practice points**

1. This systematic review examined the association between sleep duration and mood in 361,505 adolescents from 74 identified studies, 73 of which were suitable for meta-analysis.
2. Pooled results indicated that shorter sleep was associated with a 55% increased risk of mood deficits across mood states.
3. The effect that short sleep had on mood varied according to which mood state was assessed, the study design, the operationalisation of sleep and geographical region.
4. Short sleep has a significant deleterious effect on a range of positive and negative mood states among healthy, non-clinical samples of adolescents. It is imperative that greater focus is given to sleep as for prevention and early intervention for mood deficits.

### **Research agenda**

1. While positive affect showed the largest impact from short sleep, research has largely focussed on negative affect and mood states. Future research would profit by including a range of positive mood states.
2. Dose-response studies of the effect of sleep on a range of mood states are needed to determine the degree of sleep loss at which mood deficits occur, whether they accumulate over time, and the optimal duration of sleep needed for mood.
3. Disorder-specific models of psychiatric conditions include valuable information on those aspects of affective and cognitive-affective functioning that precipitate and maintain mental ill-health. Research is needed that examines mood from a much broader framework to test whether sleep affects mental health through these disorder-specific mechanisms.
4. Research would benefit by including broader aspects of sleep in addition to sleep duration, such as sleep quality, sleep timing, sleep efficiency, sleep regularity and chronotype, as they may all have a unique impact on mood.



## REFERENCES

1. Thelwell RC, Lane AM, Weston NJ. Mood states, self-set goals, self-efficacy and performance in academic examinations. *Personality and Individual Differences*. 2007;42(3):573-83.
2. Jacobson KC, Rowe DC. Genetic and environmental influences on the relationships between family connectedness, school connectedness, and adolescent depressed mood: sex differences. *Dev Psychol*. 1999;35(4):926-39.
3. Butler JM, Whalen CK, Jamner LD. Bummed out now, feeling sick later: weekday versus weekend negative affect and physical symptom reports in high school freshmen. *The Journal of adolescent health : official publication of the Society for Adolescent Medicine*. 2009;44(5):452-7.
4. Paus T, Keshavan M, Giedd JN. Why do many psychiatric disorders emerge during adolescence? *Nat Rev Neurosci*. 2008;9(12):947-57.
5. Merikangas KR, He JP, Burstein M, Swanson SA, Avenevoli S, Cui L, et al. Lifetime prevalence of mental disorders in U.S. adolescents: results from the National Comorbidity Survey Replication--Adolescent Supplement (NCS-A). *Journal of the American Academy of Child and Adolescent Psychiatry*. 2010;49(10):980-9.
6. Kessler RC, Avenevoli S, Merikangas KR. Mood disorders in children and adolescents: an epidemiologic perspective. *Biological psychiatry*. 2001;49(12):1002-14.
7. Van Ameringen M, Mancini C, Farvolden P. The impact of anxiety disorders on educational achievement. *Journal of anxiety disorders*. 2003;17(5):561-71.
8. Vigo D, Thornicroft G, Atun R. Estimating the true global burden of mental illness. *The Lancet Psychiatry*. 2016;3(2):171-8.
9. Reddy MS. Depression: The Disorder and the Burden. *Indian Journal of Psychological Medicine*. 2010;32(1):1-2.
10. Gregory AM, Sadeh A. Annual Research Review: Sleep problems in childhood psychiatric disorders--a review of the latest science. *Journal of child psychology and psychiatry, and allied disciplines*. 2016;57(3):296-317.
11. Gregory AM, Sadeh A. Sleep, emotional and behavioral difficulties in children and adolescents. *Sleep Med Rev*. 2012;16(2):129-36.
12. Watling J, Pawlik B, Scott K, Booth S, Short MA. Sleep Loss and Affective Functioning: More Than Just Mood. *Behavioral sleep medicine*. 2017;15(5):394-409.
13. Kahn M, Sheppes G, Sadeh A. Sleep and emotions: bidirectional links and underlying mechanisms. *International journal of psychophysiology : official journal of the International Organization of Psychophysiology*. 2013;89(2):218-28.
- \*14. Fuligni AJ, Bai S, Krull JL, Gonzales NA. Individual Differences in Optimum Sleep for Daily Mood During Adolescence. *Journal of clinical child and adolescent psychology : the official journal for the Society of Clinical Child and Adolescent Psychology, American Psychological Association, Division 53*. 2017:1-11.
15. Shochat T, Cohen-Zion M, Tzischinsky O. Functional consequences of inadequate sleep in adolescents: a systematic review. *Sleep Med Rev*. 2014;18(1):75-87.
16. Short MA, Weber N. Sleep Duration and Risk-Taking in Adolescents: A systematic review and meta-analysis. *Sleep Medicine Reviews*. 2018.
17. Galland BC, Short MA, Terrill P, Rigney G, Haszard JJ, Coussens S, et al. Establishing normal values for pediatric nighttime sleep measured by actigraphy: a systematic review and meta-analysis. *Sleep*. 2018:zsy017-zsy.
18. Krueger PM, Friedman EM. Sleep duration in the United States: a cross-sectional population-based study. *American journal of epidemiology*. 2009;169(9):1052-63.
19. Paruthi S, Brooks LJ, D'Ambrosio C, Hall WA, Kotagal S, Lloyd RM, et al. Recommended Amount of Sleep for Pediatric Populations: A Consensus Statement of the American Academy of Sleep Medicine. *Journal of clinical sleep medicine : JCSM : official publication of the American Academy of Sleep Medicine*. 2016;12(6):785-6.

20. Short MA, Weber N, Reynolds C, Coussens S, Carskadon MA. Estimating Adolescent Sleep Need Using Dose-Response Modelling. *Sleep*. 2018;zy011-zsy.
21. Short MA, Gradisar M, Lack LC, Wright HR, Dohnt H. The sleep patterns and well-being of Australian adolescents. *Journal of adolescence*. 2013;36(1):103-10.
22. Wolfson AR, Carskadon MA. Sleep schedules and daytime functioning in adolescents. *Child development*. 1998;69(4):875-87.
23. Lovato N, Gradisar M. A meta-analysis and model of the relationship between sleep and depression in adolescents: Recommendations for future research and clinical practice. *Sleep medicine reviews*. 2014;18(6):521-9.
24. WHO. *International Statistical Classification of Diseases and Related Health Problems 10th ed*. Geneva: World Health Organisation; 1992.
25. Yu IT, Tse SL. Workshop 6--sources of bias in cross-sectional studies; summary on sources of bias for different study designs. *Hong Kong medical journal = Xianggang yi xue za zhi*. 2012;18(3):226-7.
26. Melsen WG, Bootsma MC, Rovers MM, Bonten MJ. The effects of clinical and statistical heterogeneity on the predictive values of results from meta-analyses. *Clinical microbiology and infection : the official publication of the European Society of Clinical Microbiology and Infectious Diseases*. 2014;20(2):123-9.
27. Cooper H, Hedges LV, Valentine JC. *The handbook of research synthesis and meta-analysis 2nd edition. The Hand of Res Synthesis and Meta-Analysis, 2nd Ed*: Russell Sage Foundation; 2009. p. 1-615.
28. Barnes JC, Meldrum RC. The Impact of Sleep Duration on Adolescent Development: A Genetically Informed Analysis of Identical Twin Pairs. *J Youth Adolesc*. 2014;44(2):489-506.
29. Bauducco SV, Flink IK, Jansson-Fröjmark M, Linton SJ. Sleep duration and patterns in adolescents: correlates and the role of daily stressors. *Sleep Health*. 2016;2(3):211-8.
- \*30. Baum KT, Desai A, Field J, Miller LE, Rausch J, Beebe DW. Sleep restriction worsens mood and emotion regulation in adolescents. *Journal of Child Psychology and Psychiatry*. 2014;55(2):180-90.
31. Beebe DW, Fallone G, Godiwala N, Flanigan M, Martin D, Schaffner L, et al. Feasibility and behavioral effects of an at-home multi-night sleep restriction protocol for adolescents. *Journal of Child Psychology and Psychiatry*. 2008;49(9):915-23.
32. Bei B, Byrne ML, Ivens C, Waloszek J, Woods MJ, Dudgeon P, et al. Pilot study of a mindfulness-based, multi-component, in-school group sleep intervention in adolescent girls. *Early intervention in psychiatry*. 2013;7(2):213-20.
33. Bei BDC, Manber R, Allen NB, Trinder J, Wiley JF. Too Long, Too Short, or Too Variable? Sleep Intraindividual Variability and Its Associations With Perceived Sleep Quality and Mood in Adolescents During Naturalistically Unconstrained Sleep. *Sleep*. 2017;40(2).
34. Boergers J, Gable CJ, Owens JA. Later school start time is associated with improved sleep and daytime functioning in adolescents. *Journal of Developmental and Behavioral Pediatrics*. 2014;35(1):11-7.
35. Bonnar D, Gradisar M, Moseley L, Coughlin AM, Cain N, Short MA. Evaluation of novel school-based interventions for adolescent sleep problems: Does parental involvement and bright light improve outcomes? *Sleep Health*. 2015;1(1):66-74.
36. Chan CS, Poon CYS, Leung JCY, Lau KNT, Lau EYY. Delayed school start time is associated with better sleep, daytime functioning, and life satisfaction in residential high-school students. *Journal of adolescence*. 2018;66:49-54.
37. Chue AE, Gunthert KC, Kim RW, Alfano CA, Ruggiero AR. The role of sleep in adolescents' daily stress recovery: Negative affect spillover and positive affect bounce-back effects. *Journal of adolescence*. 2018;66:101-11.

38. Conklin AI, Yao CA, Richardson CG. Chronic sleep deprivation and gender-specific risk of depression in adolescents: a prospective population-based study. *BMC public health*. 2018;18(1):724.
- \*39. Dewald-Kaufmann JF, Oort FJ, Meijer AM. The effects of sleep extension and sleep hygiene advice on sleep and depressive symptoms in adolescents: A randomized controlled trial. *Journal of Child Psychology and Psychiatry*. 2014;55(3):273-83.
40. Díaz-Morales JF. Anxiety during adolescence: Considering morningness-eveningness as a risk factor. *Sleep Biol Rhythms*. 2016;14(2):141-7.
41. Doane LD, Thurston EC. Associations among sleep, daily experiences, and loneliness in adolescence: Evidence of moderating and bidirectional pathways. *Journal of adolescence*. 2014;37(2):145-54.
42. Doane LD, Gress-Smith JL, Breitenstein RS. Multi-method assessments of sleep over the transition to college and the associations with depression and anxiety symptoms. *Journal of Youth and Adolescence*. 2015;44(2):389-404.
- \*43. El-Sheikh M, Tu KM, Saini EK, Fuller-Rowell TE, Buckhalt JA. Perceived discrimination and youths' adjustment: Sleep as a moderator. *Journal of sleep research*. 2016;25(1):70-7.
44. Fan F, Zhou Y, Liu X. Sleep Disturbance Predicts Posttraumatic Stress Disorder and Depressive Symptoms: A Cohort Study of Chinese Adolescents. *J Clin Psychiatry*. 2017;78(7):882-8.
45. Fredriksen K, Rhodes J, Reddy R, Way N. Sleepless in Chicago: Tracking the Effects of Adolescent Sleep Loss During the Middle School Years. *Child development*. 2004;75(1):84-95.
46. Hall Brown TS. The Effects of Natural Sleep Debt on Current Mood, Working Memory, and Risk-Taking Propensity 2008.
47. Hasler JC. The effect of sleep extension on academic performance, cognitive functioning and psychological distress in adolescents. 2008.
48. Hyakutake A, Kamijo T, Misawa Y, Washizuka S, Inaba Y, Tsukahara T, et al. Cross-sectional observation of the relationship of depressive symptoms with lifestyles and parents' status among Japanese junior high school students. *Environmental health and preventive medicine*. 2016;21(4):265-73.
49. Itani O, Kaneita Y, Munezawa T, Ikeda M, Osaki Y, Higuchi S, et al. Anger and Impulsivity Among Japanese Adolescents: A Nationwide Representative Survey. *The Journal of clinical psychiatry*. 2016;77(7):e860-6.
50. Josev EK, Jackson ML, Bei B, Trinder J, Harvey A, Clarke C, et al. Sleep Quality in Adolescents With Chronic Fatigue Syndrome/Myalgic Encephalomyelitis (CFS/ME). *Journal of clinical sleep medicine : JCSM : official publication of the American Academy of Sleep Medicine*. 2017;13(9):1057-66.
51. Kalak N, Gerber M, Kirov R, Mikoteit T, Pühse U, Holsboer-Trachsler E, et al. The relation of objective sleep patterns, depressive symptoms, and sleep disturbances in adolescent children and their parents: A sleep-EEG study with 47 families. *Journal of Psychiatric Research*. 2012;46(10):1374-82.
52. Kang SG, Lee YJ, Kim SJ, Lim W, Lee HJ, Park YM, et al. Weekend catch-up sleep is independently associated with suicide attempts and self-injury in Korean adolescents. *Comprehensive psychiatry*. 2014;55(2):319-25.
53. Kelly RJ, El-Sheikh M. Reciprocal relations between children's sleep and their adjustment over time. *Developmental Psychology*. 2014;50(4):1137-47.
54. Kuo SI, Updegraff KA, Zeiders KH, McHale SM, Umana-Taylor AJ, De Jesus SA. Mexican American adolescents' sleep patterns: contextual correlates and implications for health and adjustment in young adulthood. *Journal of youth and adolescence*. 2015;44(2):346-61.
55. Lazaratou H, Anagnostopoulos DC, Vlassopoulos M, Charbilas D, Rotsika V, Tsakanikos E, et al. Predictors and characteristics of anxiety among adolescent students: A Greek sample. *Psychiatriki*. 2013;24(1):27-36.

56. Lee BH, Kang SG, Choi JW, Lee YJ. The Association between Self-reported Sleep Duration and Body Mass Index among Korean Adolescents. *Journal of Korean medical science*. 2016;31(12):1996-2001.
- \*57. Lee YJ, Cho S-J, Cho IH, Kim SJ. Insufficient sleep and suicidality in adolescents. *Sleep: Journal of Sleep and Sleep Disorders Research*. 2012;35(4):455-60.
58. Lee J. Sleep duration's association with diet, physical activity, mental status, and weight among Korean high school students. *Asia Pac J Clin Nutr*. 2017;26(5):906-13.
59. Lehto JE, Uusitalo-Malmivaara L. Sleep-related factors: Associations with poor attention and depressive symptoms. *Child: Care, Health and Development*. 2014;40(3):419-25.
60. Lemola S, Brand S, Vogler N, Perkinson-Gloor N, Allemand M, Grob A. Habitual computer game playing at night is related to depressive symptoms. *Personality and Individual Differences*. 2011;51(2):117-22.
61. Lemola S, Perkinson-Gloor N, Brand S, Dewald-Kaufmann JF, Grob A. Adolescents' electronic media use at night, sleep disturbance, and depressive symptoms in the smartphone age. *Journal of Youth and Adolescence*. 2015;44(2):405-18.
62. Lin W-H, Yi C-C. Unhealthy sleep practices, conduct problems, and daytime functioning during adolescence. *Journal of Youth and Adolescence*. 2015;44(2):431-46.
63. Liu X, Zhou H. Sleep duration, insomnia and behavioral problems among Chinese adolescents. *Psychiatry Research*. 2002;111(1):75-85.
- \*64. Lo JC, Lee SM, Teo LM, Lim J, Gooley JJ, Chee MW. Neurobehavioral Impact of Successive Cycles of Sleep Restriction With and Without Naps in Adolescents. *Sleep*. 2017;40(2).
65. Lo JC, Ong JL, Leong RL, Gooley JJ, Chee MW. Cognitive performance, sleepiness, and mood in partially sleep deprived adolescents: The Need for Sleep study. *Sleep*. 2016;39(3):687-98.
66. Lovato N, Short MA, Micic G, Hiller RM, Gradisar M. An investigation of the longitudinal relationship between sleep and depressed mood in developing teens. *Nature and science of sleep*. 2017;9:3-10.
67. Maume DJ. Social relationships and the sleep-health nexus in adolescence: evidence from a comprehensive model with bi-directional effects. *Sleep Health*. 2017;3(4):284-9.
68. Mazzer K, Boersma K, Linton SJ. A longitudinal view of rumination, poor sleep and psychological distress in adolescents. 2018;245:686-96.
- \*69. McGlinchey EL, Talbot LS, Chang K-h, Kaplan KA, Dahl RE, Harvey AG. The effect of sleep deprivation on vocal expression of emotion in adolescents and adults. *Sleep: Journal of Sleep and Sleep Disorders Research*. 2011;34(9):1233-41.
70. McMakin DL, Dahl RE, Buysse DJ, Cousins JC, Forbes EE, Silk JS, et al. The impact of experimental sleep restriction on affective functioning in social and nonsocial contexts among adolescents. *Journal of Child Psychology and Psychiatry*. 2016;57(9):1027-37.
71. Moore M, Kirchner L, Drotar D, Johnson N, Rosen C, Ancoli-Israel S, et al. Relationships among sleepiness, sleep time, and psychological functioning in adolescents. *Journal of Pediatric Psychology*. 2009;34(10):1175-83.
72. Mullin BC, Pyle L, Haraden D, Riederer J, Brim N, Kaplan D, et al. A Preliminary Multimethod Comparison of Sleep Among Adolescents With and Without Generalized Anxiety Disorder. *Journal of clinical child and adolescent psychology : the official journal for the Society of Clinical Child and Adolescent Psychology, American Psychological Association, Division 53*. 2017;46(2):198-210.
73. Norell-Clarke A, Hagquist C. Child and adolescent sleep duration recommendations in relation to psychological and somatic complaints based on data between 1985 and 2013 from 11 to 15-year-olds. *Journal of adolescence*. 2018;68:12-21.
74. Nowakowski S, Choi H, Meers J, Temple JR. Inadequate Sleep as a Mediating Variable between Exposure to Interparental Violence and Depression Severity in Adolescents. *Journal of child & adolescent trauma*. 2016;9(2):109-14.

75. Nuutinen T, Roos E, Ray C, Villberg J, Välimaa R, Rasmussen M, et al. Computer use, sleep duration and health symptoms: A cross-sectional study of 15-year olds in three countries. *International Journal of Public Health*. 2014;59(4):619-28.
76. Oginska H, Pokorski J. Fatigue and mood correlates of sleep length in three age-social groups: School children, students, and employees. *Chronobiology International*. 2006;23(6):1317-28.
- \*77. Owens JA, Belon K, Moss P. Impact of delaying school start time on adolescent sleep, mood, and behavior. *Archives of pediatrics & adolescent medicine*. 2010;164(7):608-14.
78. Pallesen S, Saxvig IW, Molde H, Sørensen E, Wilhelmsen-Langeland A, Bjorvatn B. Brief report: Behaviorally induced insufficient sleep syndrome in older adolescents: Prevalence and correlates. *Journal of adolescence*. 2010.
79. Pasch KE, Laska MN, Lytle LA, Moe SG. Adolescent sleep, risk behaviors, and depressive symptoms: Are they linked? *American Journal of Health Behavior*. 2010;34(2):237-48.
80. Randler C, Weber V. Positive and negative affect during the school day and its relationship to morningness-eveningness. *Biological Rhythm Research*. 2015;46(5):683-90.
81. Raniti MB, Allen NB, Schwartz O, Waloszek JM, Byrne ML, Woods MJ, et al. Sleep Duration and Sleep Quality: Associations With Depressive Symptoms Across Adolescence. *Behavioral sleep medicine*. 2017;15(3):198-215.
- \*82. Reddy R, Palmer CA, Jackson C, Farris SG, Alfano CA. Impact of sleep restriction versus idealized sleep on emotional experience, reactivity and regulation in healthy adolescents. *Journal of sleep research*. 2017;26(4):516-25.
83. Roberts RE, Duong HT. The prospective association between sleep deprivation and depression among adolescents. *Sleep: Journal of Sleep and Sleep Disorders Research*. 2014;37(2):239-44.
84. Sarchiapone M, Mandelli L, Carli V, Iosue M, Wasserman C, Hadlaczky G, et al. Hours of sleep in adolescents and its association with anxiety, emotional concerns, and suicidal ideation. *Sleep Medicine*. 2014;15(2):248-54.
85. Seo JH, Kim JH, Yang KI, Hong SB. Late use of electronic media and its association with sleep, depression, and suicidality among Korean adolescents. *Sleep medicine*. 2017;29:76-80.
- \*86. Shen L, van Schie J, Ditchburn G, Brook L, Bei B. Positive and Negative Emotions: Differential Associations with Sleep Duration and Quality in Adolescents. *Journal of youth and adolescence*. 2018:1-12.
87. Short MA, Louca M. Sleep deprivation leads to mood deficits in healthy adolescents. *Sleep Medicine*. 2015;16(8):987-93.
88. Short MA, Gradisar M, Lack LC, Wright HR. The impact of sleep on adolescent depressed mood, alertness and academic performance. *Journal of adolescence*. 2013;36(6):1025-33.
89. Singh R, Suri JC, Sharma R, Suri T, Adhikari T. Sleep Pattern of Adolescents in a School in Delhi, India: Impact on their Mood and Academic Performance. *Indian J Pediatr*. 2018;85(10):841-8.
90. Stheneur C, Sznajder M, Spiry C, Marcu Marin M, Ghout I, Samb P, et al. Sleep duration, quality of life and depression in adolescents: a school-based survey. 2017.
91. Suzuki H, Kaneita Y, Osaki Y, Minowa M, Kanda H, Suzuki K, et al. Clarification of the factor structure of the 12-item General Health Questionnaire among Japanese adolescents and associated sleep status. *Psychiatry Research*. 2011;188(1):138-46.
92. Van Dyk TR, Zhang N, Catlin PA, Cornist K, McAlister S, Whitacre C, et al. Feasibility and Emotional Impact of Experimentally Extending Sleep in Short-Sleeping Adolescents. *Sleep*. 2017;40(9).
93. Wahlstrom KL, Berger AT, Widome R. Relationships between school start time, sleep duration, and adolescent behaviors. *Sleep Health*. 2017.
94. Warner S, Murray G, Meyer D. Holiday and school-term sleep patterns of Australian adolescents. *Journal of adolescence*. 2008;31(5):595-608.
95. Watson A, Brickson S. Impaired Sleep Mediates the Negative Effects of Training Load on Subjective Well-Being in Female Youth Athletes. *Sports Health*. 2018;10(3):244-9.

96. Yang SJ, Cha HS. Retrospective cohort study on Korean adolescents' sleep, depression, school adjustment, and life satisfaction. 2018;20(4):422-30.
97. Yip T. The effects of ethnic/racial discrimination and sleep quality on depressive symptoms and self-esteem trajectories among diverse adolescents. *Journal of Youth and Adolescence*. 2015;44(2):419-30.
98. Hirshkowitz M, Whiton K, Albert S, Alessi C, Bruni O, DonCarlos L, et al. National Sleep Foundation's sleep time duration recommendations: methodology and results summary. *Sleep Health*. 2015;1(1):40-3.
99. Kouros CD, El-Sheikh M. Daily mood and sleep: reciprocal relations and links with adjustment problems. *J Sleep Res*. 2015;24(1):24-31.
100. Yoo SS, Gujar N, Hu P, Jolesz FA, Walker MP. The human emotional brain without sleep--a prefrontal amygdala disconnect. *Current biology : CB*. 2007;17(20):R877-8.
101. Ong JL, Lo JC, Gooley JJ, Chee MW. EEG changes across multiple nights of sleep restriction and recovery in adolescents: the need for sleep study. 2016.
102. Palmer CA, Alfano CA. Sleep and emotion regulation: An organizing, integrative review. *Sleep medicine reviews*. 2017;31:6-16.
103. Nota JA, Coles ME. Shorter sleep duration and longer sleep onset latency are related to difficulty disengaging attention from negative emotional images in individuals with elevated transdiagnostic repetitive negative thinking. *Journal of behavior therapy and experimental psychiatry*. 2018;58:114-22.
104. Short MA, Blunden S, Rigney G, Matricciani L, Coussens S, M. Reynolds C, et al. Cognition and objectively measured sleep duration in children: a systematic review and meta-analysis. *Sleep Health*. 2018;4(3):292-300.

## Appendix A

## Literature Search

Source	Search string	Result	Notes
<b>Source:</b> PubMed  <b>Coverage:</b> 1809-  <b>Search Date:</b> 2018-12-17	(("Adolescent"[Mesh] OR adolescen*[Title/Abstract] OR youth*[Title/Abstract] OR teenager*[Title/Abstract] OR teen[Title/Abstract] OR teens[Title/Abstract]) AND ("sleep duration"[Title/Abstract] OR "sleep deprivation"[Title/Abstract] OR "Sleep Deprivation"[Mesh] OR "sleep loss"[Title/Abstract] OR "total sleep"[Title/Abstract] OR "sleep restriction"[Title/Abstract]) AND ("Confusion"[Mesh] OR happiness[Title/Abstract] OR suicidal[Title/Abstract] OR "Suicidal Ideation"[Mesh] OR antagonism[Title/Abstract] OR apathy[Title/Abstract] OR "Apathy"[Mesh] OR oppositionality [Title/Abstract] OR euphori*[Title/Abstract] OR "Euphoria"[Mesh] OR "Depression"[Mesh] OR depress*[Title/Abstract] OR tense*[Title/Abstract] OR tension[Title/Abstract] OR irrita*[Title/Abstract] OR annoyed[Title/Abstract] OR hostil*[Title/Abstract] OR "Hate"[Mesh] OR "Happiness"[Mesh] OR hate*[Title/Abstract] OR "Hostility"[Mesh] OR frustrat*[Title/Abstract] OR "Frustration"[Mesh] OR "Anxiety"[Mesh] OR rage[Title/Abstract] OR "Anger"[Mesh] OR "Rage"[Mesh] OR "Irritable Mood"[Mesh] OR moods[Title/Abstract] OR mood[Title/Abstract] OR moody[Title/Abstract] OR anxious*[Title/Abstract] OR emotion*[Title/Abstract] OR temper[Title/Abstract] OR anxiety*[Title/Abstract] OR distress*[Title/Abstract] OR anger*[Title/Abstract] OR angry[Title/Abstract] OR happy*[Title/Abstract] OR sad[Title/Abstract] OR sadness[Title/Abstract] OR confus*[Title/Abstract] OR upset[Title/Abstract]) AND (English[lang]))	<b>781</b>	All terms searched in the fields "Title/Abstract" and in "MeSH" when available  <b>Filters applied:</b> English Language

<p><b>Source:</b> Scopus</p> <p><b>Coverage:</b> 1996-</p> <p><b>Search Date:</b> 2018-12-14</p>	<p>(( TITLE ( "sleep duration" OR "sleep deprivation" OR "sleep loss" OR "total sleep" OR "sleep restriction" ) OR ABS ( "sleep duration" OR "sleep deprivation" OR "sleep loss" OR "total sleep" OR "sleep restriction" ) ) ) AND ( ( TITLE ( adolescen* OR youth* OR teenager* OR teen OR teens ) OR ABS ( adolescen* OR youth* OR teenager* OR teen OR teens ) ) ) AND ( ( TITLE ( happiness OR suicidal OR antagonism OR apathy OR oppositionality OR euphori* OR depress* OR tense* OR tension OR irrita* OR annoyed OR hostile* OR hate* OR frustrat* OR rage OR moods OR mood OR moody ) OR ABS ( happiness OR suicidal OR antagonism OR apathy OR oppositionality OR euphori* OR depress* OR tense* OR tension OR irrita* OR annoyed OR hostile* OR hate* OR frustrat* OR rage OR moods OR mood OR moody ) OR ABS ( anxious* OR emotion* OR temper OR anxiety* OR distress* OR anger* OR angry OR happy* OR sad OR sadness OR confus* OR upset ) OR TITLE ( anxious* OR emotion* OR temper OR anxiety* OR distress* OR anger* OR angry OR happy* OR sad OR sadness OR confus* OR upset ) ) ) AND ( LIMIT-TO ( LANGUAGE , "English" ) )</p>	<p><b>344</b></p>	<p>All terms searched in the fields "TITLE" and "ABS" (Abstract). No thesaurus available.</p> <p><b>Filters applied:</b> English Language</p>
<p><b>Source:</b> PsychInfo (EBSCO)</p> <p><b>Coverage:</b> 1987-</p>	<p>((TI ( happiness OR suicidal OR antagonism OR apathy OR oppositionality OR euphori* OR depress* OR tense* OR tension OR irrita* OR annoyed OR hostile* OR hate* OR frustrat* OR rage OR moods OR mood OR moody OR anxious* OR temper OR emotion* OR anxiety OR distress* OR anger* OR angry OR happy* OR sad OR sadness OR confus* OR upset ) OR AB ( happiness OR suicidal OR antagonism OR apathy OR oppositionality OR euphori* OR depress* OR tense* OR tension OR irrita* OR annoyed OR hostile* OR hate* OR frustrat* OR rage OR moods OR mood OR moody OR anxious* OR temper OR emotion* OR anxiety OR distress* OR anger* OR</p>	<p><b>292</b></p>	<p>All terms searched in the fields "Title" (TI) and "Abstract" (AB) and in the thesaurus (DE) when available.</p> <p><b>Filters applied:</b> English Language</p>



<p><b>Search Date:</b> 2018-12-14</p>	<p>angry OR happy* OR sad OR sadness OR confus* OR upset ) OR (DE ("Mental Confusion" OR "Happiness" OR "Suicidal Ideation" OR "Hostility" OR "Apathy" OR "Euphoria" OR "Major Depression" OR "Depression Emotion" OR "Irritability" OR "Hate" OR "Frustration" OR "Anger" OR "Moodiness" OR "anxiety" OR "distress" OR "sadness")</p> <p>AND (TI ( "sleep duration" OR "sleep deprivation" OR "sleep loss" OR "total sleep" OR "sleep restriction" ) OR AB ( "sleep duration" OR "sleep deprivation" OR "sleep loss" OR "total sleep" OR "sleep restriction" ) OR DE ("Sleep Deprivation") AND (TI ( adolescen* OR youth* OR teenager* OR teen OR teens ) OR AB ( adolescen* OR youth* OR teenager* OR teen OR teens ))</p>		
<p><b>Source:</b> Embase (Elsevier)</p> <p><b>Coverage:</b> 1947-</p> <p><b>All sources included</b></p> <p><b>Search Date:</b></p>	<p>((('sleep duration':ab,ti OR 'sleep deprivation':ab,ti OR 'sleep loss':ab,ti OR 'total sleep':ab,ti OR 'sleep restriction':ab,ti OR 'sleep time'/de OR 'sleep deprivation'/de) AND (adolescen*:ab,ti OR youth*:ab,ti OR teenager*:ab,ti OR teen:ab,ti OR teens:ab,ti OR 'adolescence'/de) AND (happiness:ab,ti OR suicidal:ab,ti OR antagonism:ab,ti OR apathy:ab,ti OR oppositionality:ab,ti OR euphori*:ab,ti OR depress*:ab,ti OR tense*:ab,ti OR tension:ab,ti OR irrita*:ab,ti OR annoyed:ab,ti OR hostil*:ab,ti OR hate*:ab,ti OR frustrat*:ab,ti OR rage:ab,ti OR moods:ab,ti OR mood:ab,ti OR moody:ab,ti OR anxious*:ab,ti OR temper:ab,ti OR emotion*:ab,ti OR anxiety:ab,ti OR distress*:ab,ti OR anger*:ab,ti OR angry:ab,ti OR happy*:ab,ti OR sad:ab,ti OR sadness:ab,ti OR confus*:ab,ti OR upset:ab,ti OR 'suicidal ideation'/de OR 'apathy'/de OR 'euphoria'/de OR 'depression'/de OR 'irritability'/de OR 'hostility'/de OR 'hate'/de OR 'frustration'/de OR 'sadness'/de OR 'happiness'/de OR 'anger'/de OR 'anxiety'/de OR 'mood'/de OR 'rage'/de OR 'confusion'/de) AND</p>	<p><b>745</b></p>	<p>All terms searched in the fields Title” and “Abstract” (ab,ti) and in “MeSH” (/de) when available</p> <p><b>Filters applied:</b> English Language</p>

2018-12-14	([english]/lim))		
<b>Total no. of references located</b>		<b>2162</b>	
<b>Total no. of references after de- duplication</b>		<b>1532</b>	





Figure 1. PRISMA diagram showing the results of the literature search.

Journal Pre-proof






Figure 2. Forest plot showing effect sizes for individual studies, together with the random overall effect size in odds ratios.




Figure 3. Risk of bias assessment. A study was deemed as having partial risk of bias for an indicator if information was not provided regarding that indicator.




Figure 4. Funnel plot of effect sizes and their standard error.

Journal Pre-proof