

2-1-2022

Starlike Functions of Complex Order with Respect to Symmetric Points Defined Using Higher Order Derivatives

Kadhavoor R. Karthikeyan

National University of Science & Technology (by Merger of Caledonian College of Engineering and Oman Medical College)

Sakkarai Lakshmi

University of Technology and Applied Sciences–Al Musannah

Seetharam Varadharajan

University of Technology and Applied Sciences–Al Musannah

Dharmaraj Mohankumar

University of Madras

Elangho Umadevi

Zayed University

Follow this and additional works at: <https://zuscholars.zu.ac.ae/works>



Part of the [Mathematics Commons](#)

Recommended Citation

Karthikeyan, Kadhavoor R.; Lakshmi, Sakkarai; Varadharajan, Seetharam; Mohankumar, Dharmaraj; and Umadevi, Elangho, "Starlike Functions of Complex Order with Respect to Symmetric Points Defined Using Higher Order Derivatives" (2022). *All Works*. 4897.

<https://zuscholars.zu.ac.ae/works/4897>

This Article is brought to you for free and open access by ZU Scholars. It has been accepted for inclusion in All Works by an authorized administrator of ZU Scholars. For more information, please contact scholars@zu.ac.ae.



Article

Starlike Functions of Complex Order with Respect to Symmetric Points Defined Using Higher Order Derivatives

Kadhavoor R. Karthikeyan ^{1,*} , Sakkarai Lakshmi ² , Seetharam Varadharajan ² , Dharmaraj Mohankumar ³ and Elangho Umadevi ⁴ ¹ Department of Applied Mathematics and Science, National University of Science & Technology, Muscat P.O. Box 620, Oman² Mathematics Section, Department of Information Technology, University of Technology and Applied Sciences–Al Musannah, Musannah P.O. Box 191, Oman; laxmirmk@gmail.com (S.L.); varadharajan@act.edu.om (S.V.)³ P.G. and Research Department of Mathematics, Pachaiyappa’s College, University of Madras, Chennai 600030, India; dmohankumarmaths@gmail.com⁴ Department of Mathematics and Statistics, College of Natural and Health Sciences, Zayed University, Abu Dhabi P.O. Box 144534, United Arab Emirates; z10011@zu.ac.ae

* Correspondence: karthikeyan@nu.edu.om; Tel.: +968-95159288

Abstract: In this paper, we introduce and study a new subclass of multivalent functions with respect to symmetric points involving higher order derivatives. In order to unify and extend various well-known results, we have defined the class subordinate to a conic region impacted by Janowski functions. We focused on conic regions when it pertained to applications of our main results. Inclusion results, subordination property and coefficient inequality of the defined class are the main results of this paper. The applications of our results which are extensions of those given in earlier works are presented here as corollaries.

Keywords: multivalent functions; starlike and convex functions; coefficient inequalities; analytic function; univalent function; Schwartz function; differential subordination; Fekete-Szegő inequality



Citation: Karthikeyan, K.R.; Lakshmi, S.; Varadharajan, S.; Mohankumar, D.; Umadevi, E. Starlike Functions of Complex Order with Respect to Symmetric Points Defined Using Higher Order Derivatives. *Fractal Fract.* **2022**, *6*, 116. <https://doi.org/10.3390/fractalfract6020116>

Academic Editors: Acu Mugur Alexandru and Shahram Najafzadeh

Received: 27 January 2022

Accepted: 7 February 2022

Published: 17 February 2022

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction and Definitions

Throughout this paper, we let \mathbb{C} , \mathbb{Z}^- and \mathbb{N} to denote the sets of complex numbers, negative integers and natural numbers, respectively. Let $\mathcal{H}(a, n)$ be the class comprising of all analytic functions defined in unit disc $\mathbb{E} = \{z \in \mathbb{C} : |z| < 1\}$ and having a power series representation of the form $h(z) = a + a_n z^n + a_{n+1} z^{n+1} + \dots$. Furthermore, let $\mathcal{A}(p, n)$ denote the class of functions h analytic in \mathbb{E} and having a power series representation of the form

$$h(z) = z^p + \sum_{k=p+n}^{\infty} a_k z^k, \quad (p, n \in \mathbb{N}) \quad (1)$$

and let $\mathcal{A}(1, 1) = \mathcal{A}$. Two prominent subclasses of \mathcal{A} are the so-called families of starlike functions and convex functions which have the analytic characterization of the form

$$\operatorname{Re}\left(\frac{zh'(z)}{h(z)}\right) > 0 \quad \text{and} \quad \operatorname{Re}\left(1 + \frac{zh''(z)}{h'(z)}\right) > 0,$$

respectively. Here we let \mathcal{S}^* and \mathcal{C} to denote the class of starlike functions and convex functions, respectively. The two preceding descriptions reveal an interesting close analytic characterization between starlike and convex functions. This says that $h(z) \in \mathcal{C}$ if and only if $zh'(z) \in \mathcal{S}^*$. For detailed study and developments pertaining to various subclasses of $\mathcal{A}(p, n)$, refer to [1,2]. We let the collection \mathcal{P} of functions $\psi(z)$ that are analytic in the unit

disc \mathbb{E} with $\psi(0) = 1$ and $\operatorname{Re} \psi(z) > 0$. Hereafter, we let $\psi \in \mathcal{P}$ and ψ , has a power series expansion of the form

$$\psi(z) = 1 + L_1z + L_2z^2 + L_3z^3 + \dots, \quad z \in \mathbb{E}, \quad L_1 > 0. \quad (2)$$

Subordination, quasi-subordination and Hadamard product (or convolution) are the three main tools that are predominantly used in the study of univalent functions theory. We let \prec , \prec_{κ} and $*$ to denote the subordination, quasi-subordination and Hadamard product, respectively. For detailed discussion and formal definition of the quasi-subordination and Hadamard product, refer to [3,4].

Using the principal of subordination, Ma and Minda [5] defined the classes $\mathcal{S}^*(\psi)$ and $\mathcal{C}(\psi)$ as follows:

$$\mathcal{S}^*(\psi) = \left\{ h \in \mathcal{A} : \frac{zh'(z)}{h(z)} \prec \psi(z) \right\} \quad \text{and} \quad \mathcal{C}(\psi) = \left\{ h \in \mathcal{A} : 1 + \frac{zh''(z)}{h'(z)} \prec \psi(z) \right\},$$

where $\psi(z)$ is defined as in (2). They assumed the superordinate function ψ maps the open unit disc \mathbb{E} onto a starlike region with respect to 1 and symmetric with respect to the real axis. The classes $\mathcal{S}^*(\psi)$ and $\mathcal{C}(\psi)$ consolidated the study of several generalizations of starlike and convex functions. By restricting the ψ to a specific conic region for example to parabola, cardioid and Bernoulli lemniscate, several authors studied the properties of starlike functions with respect to conic regions. Most popular among the study of starlike functions associated with conic regions are the classes $\mathcal{S}^*(\sqrt{1+z})$ defined by Sokół [6] and followed by $\mathcal{S}^*(z + \sqrt{1+z^2})$ defined by Raina and Sokół [7]. For studies related to conic region, refer to [8–15] and references provided therein.

The famous Janowski starlike functions and Janowski convex functions (see [16]), are denoted by the special case of $\mathcal{S}^*(\psi)$ and $\mathcal{C}(\psi)$, although they are still in spotlight due to their versatility. We denote by $\mathcal{S}^*(F, G)$ and $\mathcal{C}(F, G)$ the class of Janowski starlike functions and Janowski convex functions, defined by

$$\mathcal{S}^*(F, G) := \left\{ h \in \mathcal{A} : \frac{zh'(z)}{h(z)} \prec \frac{1+Fz}{1+Gz}, -1 \leq G < F \leq 1 \right\},$$

and

$$\mathcal{C}(F, G) := \left\{ h \in \mathcal{A} : 1 + \frac{zh''(z)}{h'(z)} \prec \frac{1+Fz}{1+Gz}, -1 \leq G < F \leq 1 \right\},$$

respectively. It should be noted that all the classes mentioned above were extended for $h(z)$, which belongs to $\mathcal{A}(p, 1)$. Extending the well-known Janowski class of functions [16], Aouf [17] (Equation (1.4)) defined the class $\ell(z) \in \mathcal{P}(F, G, p, \alpha)$ if and only if

$$\ell(z) = \frac{p + [pG + (F - G)(p - \alpha)]w(z)}{[1 + Gw(z)]}, \quad (-1 \leq G < F \leq 1, 0 \leq \alpha < 1) \quad (3)$$

for all $z \in \mathbb{E} = \{z : |z| < 1\}$ where $w(z)$ is the Schwartz function. Recently, Breaz et al. [18] (Equation (4)) used the following expression to study a new class of multivalent function

$$\aleph(p; F, G; \alpha; \psi; z) = \frac{[(1+F)p + \alpha(G-F)]\psi(z) + [(1-F)p - \alpha(G-F)]}{[(G+1)\psi(z) + (1-G)]}, \quad (4)$$

where $\psi(z)$ is defined as in (2). $\aleph(p; F, G; \alpha; \psi; z)$ is an extension of the class $\mathcal{P}(F, G, p, \alpha)$. Refer to [18,19], for an explanation of the purpose and motivation in order to define a class of functions superordinate to $\aleph(p; F, G; \alpha; \psi; z)$.

Recently, Aouf, Bulboacă and Seoudy in [20] (Definition 1) introduced a class so-called multivalent non-Bazilevič functions as follows: A function $h \in \mathcal{A}(p, n)$ is said to be in $\mathcal{N}_p^n(\lambda, \beta, m; F, G)$ if it satisfies

$$(1 + \lambda) \left(\frac{\delta(p, m)z^{p-m}}{h^{(m)}(z)} \right)^\beta - \lambda \frac{zh^{(1+m)}(z)}{(p-m)h^{(m)}(z)} \left(\frac{\delta(p, m)z^{p-m}}{h^{(m)}(z)} \right)^\beta \prec \frac{1 + Fz}{1 + Gz}$$

$$(\lambda \in \mathbb{C}, 0 < \beta < 1, p, n \in \mathbb{N}, m \in \mathbb{N}_0 = \mathbb{N} \cup \{0\}, p > m \text{ and } -1 \leq G < F \leq 1),$$

where \prec denotes usual subordination of analytic functions and $\delta(p, m) = \frac{p!}{(p-m)!}$.

Using Hadamard product (or convolution), Karthikeyan et al. [21] (Definition 1.1) defined a class $\mathcal{PS}_\delta^\lambda(\beta, \theta; b; \psi; h; F, G)$ of $\mathcal{A}(1, 1)$ subject to satisfying the condition

$$1 + \frac{(1 + i \tan \theta)}{b} \left[\frac{z^{1-\lambda} [\mathcal{R}'(z)]^\delta}{[(1-\beta)\mathcal{R}(z) + \beta z]^{1-\lambda}} - 1 \right] \prec \aleph(1; F, G; 0; \psi; z),$$

where $\mathcal{R} = h * g, -\frac{\pi}{2} < \theta < \frac{\pi}{2}, \delta \geq 1, 0 \leq \beta \leq 1, \lambda \geq 0, b \in \mathbb{C} \setminus \{0\}$ and $\aleph(p; F, G; \alpha; \psi; z)$ is defined as in (4).

1.1. Motivation, Novelty and Discussion

Motivated by the classes $\mathcal{N}_p^n(\lambda, \beta, m; F, G)$ and $\mathcal{PS}_\delta^\lambda(\beta, \theta; b; \psi; h; F, G)$, we aim to define and study an interesting subclass of multivalent functions with respect to symmetric points subordinate to $\aleph(p; F, G; \alpha; \psi; z)$. However, the present study is not a direct generalization or unification of $\mathcal{N}_p^n(\lambda, \beta, m; F, G)$ and $\mathcal{PS}_\delta^\lambda(\beta, \theta; b; \psi; h; F, G)$, but is closely related to the above defined function classes.

This paper is structured as follows. In this section, we will begin by illustrating that impact of $\aleph(p; F, G; \alpha; \psi; z)$ is not same on all conic regions and it varies from region to region. Subsequently, we define a class of multivalent functions using higher order derivatives superordinated by $\aleph(p; F, G; \alpha; \psi; z)$. In the Section 2, we discuss some elementary and known results which would be used to obtain our main results. Sections 3 and 4 are devoted to provide our main results namely solution to the Fekete-Szegő problem and interesting subordination conditions. Finally attempting the discretization of our results, we study the same defined function class by replacing the ordinary derivative with q -difference operator.

In [18], the geometrical interpretation and the impact of $\aleph(p; F, G; \alpha; \psi; z)$ on various conic region was not discussed in detail. Here we will consider few conic regions and we will illustrate the impact of $\aleph(p; F, G; \alpha; \psi; z)$ on $\psi(z)$. For uniformity, the colour of graphs have been based on the parameter values, which are as follows: *Red colour* is used when $\aleph(1; 1, -1; 0; \psi; z)$; *Blue colour* is used if $\aleph(1; 0, -0.3; 0.9; \psi; z)$; *Green colour* is used if $\aleph(1; 0.9, 0.0; 0.5; \psi; z)$; and *Yellow colour* is used if $\aleph(1; 0.9, 0.8; 0.8; \psi; z)$.

1.2. Comparison on The Impact of $\aleph(p; F, G; \alpha; \psi; z)$ on Two Different Conic Regions

The behaviour or impact of $\aleph(p; F, G; \alpha; \psi; z)$ is not same on all conic region ψ . To illustrate this fact, we consider two functions which maps unit disc on to a conic region of same shape namely

1. Cardioid region with cusp on the right hand side, $(\psi(z) = \frac{3+2z-z^2}{2})$.
2. Cardioid region with cusp on left hand side, $(\psi(z) = 1 + \frac{z}{k} \left(\frac{k+z}{k-z} \right), k = 1 + \sqrt{2})$.

We begin the illustration with the following.

1. It is well-known that $\psi(z) = \frac{3+2z-z^2}{2}$ is univalent in \mathbb{E} and maps the unit disc onto the interior of the cardioid with cusp on the right hand side in the right half plane (see Figure 1a). Note that while $\text{Re}[\psi(z)] = \text{Re} \left[\frac{3+2z-z^2}{2} \right] > 0$, it does not have the usual normalization $\psi(0) = 1$. The impact of $\aleph(p; F, G; \alpha; \psi; z)$ on $\psi(z) = \frac{3+2z-z^2}{2}$ is that the map is circular if F and G are chosen remotely (far off), while the curves are

polygonal (see Figure 1d) if F and G are chosen close enough. The presence of α is helpful in translation.




Figure 1. Impact of $\aleph(p; F, G; \alpha; \psi; z)$ on the conic region $\psi(z) = \frac{3}{2} + z - \frac{z^2}{2}$. (a) Mapping of \mathbb{E} under the transformation $\psi(z) = \frac{3}{2} + z - \frac{z^2}{2}$. (b) Mapping of \mathbb{E} under the transformation $\aleph(1; 0, -0.3; 0.9; \psi; z)$ if $\psi(z) = \frac{3}{2} + z - \frac{z^2}{2}$. (c) Mapping of \mathbb{E} under the transformation $\aleph(1; 0.9, 0.0; 0.5; \psi; z)$ if $\psi(z) = \frac{3}{2} + z - \frac{z^2}{2}$. (d) Mapping of \mathbb{E} under the transformation $\aleph(1; 0.9, 0.8; 0.8; \psi; z)$ if $\psi(z) = \frac{3}{2} + z - \frac{z^2}{2}$.

2. Now, if we choose

$$\begin{aligned} \psi(z) &= 1 + \frac{z}{k} \left(\frac{k+z}{k-z} \right), \quad (k = 1 + \sqrt{2}), \\ &= 1 + \frac{z}{1 + \sqrt{2}} + \frac{2z^2}{(1 + \sqrt{2})^2} + \cdots + \frac{2z^n}{(1 + \sqrt{2})^n} + O[z]^{n+1}. \end{aligned}$$

We can easily see that the function has a normalization $\psi(0) = 1$, $\text{Re}[\psi(z)] > 0$ and maps unit disc on to the cardioid with cusp on the left hand side (see Ahuja et al. [22]). From Figure 2a–d, we find that there is no major changes to the conic.




Figure 2. Impact of $\aleph(p; F, G; \alpha; \psi; z)$ on the conic region $\psi(z) = 1 + \frac{z}{k} \left(\frac{k+z}{k-z} \right)$. (a) Mapping of \mathbb{E} under the transformation $\psi(z) = 1 + \frac{z}{k} \left(\frac{k+z}{k-z} \right)$. (b) Mapping of \mathbb{E} under the transformation $\aleph(1; 0, -0.3; 0.9; \psi; z)$ if $\psi(z) = 1 + \frac{z}{k} \left(\frac{k+z}{k-z} \right)$. (c) Mapping of \mathbb{E} under the transformation $\aleph(1; 0.9, 0.0; 0.5; \psi; z)$ if $\psi(z) = 1 + \frac{z}{k} \left(\frac{k+z}{k-z} \right)$. (d) Mapping of \mathbb{E} under the transformation $\aleph(1; 0.9, 0.8; 0.8; \psi; z)$ if $\psi(z) = 1 + \frac{z}{k} \left(\frac{k+z}{k-z} \right)$.

Notice that Figures 1 and 2 have been assigned same set parameter values, the only difference being different $\psi(z)$. Comparing Figures 1 and 2, we see that the behaviour of $\aleph(p; F, G; \alpha; \psi; z)$ on various conic regions are not same (also see Noor and Malik [19]).

If $\psi \in \mathcal{P}$, then $\aleph(p; F, G; \alpha; \psi; 0) = p$ and $\text{Re}(\aleph(p; F, G; \alpha; \psi; z)) > 0$. We say that $\aleph(p; F, G; \alpha; \psi; z) \in \mathcal{P}(F, G, p, \alpha)$ if and only if it satisfies (3). We denote by $\mathcal{S}_p^*(F, G; \alpha; \psi)$ and $\mathcal{C}_p(F, G; \alpha; \psi)$, the classes of functions satisfying the condition $\frac{zh'(z)}{h(z)} \prec \aleph(p; F, G; \alpha; \psi; z)$ and $1 + \frac{zh''(z)}{h'(z)} \prec \aleph(p; F, G; \alpha; \psi; z)$, respectively. Additionally, $\mathcal{S}_1^*(1, -1; 0; \psi) := \mathcal{S}^*(\psi)$ and $\mathcal{C}_1(1, -1; 0; \psi) := \mathcal{C}(\psi)$.

The function $p_{\nu,\sigma}(\zeta)$, that plays the role of an extremal function related to the conic domain, is given by

$$\hat{p}_{\nu,\sigma}(z) = \begin{cases} \frac{1+(1-2\sigma)z}{1-z}, & \text{if } \nu = 0 \\ 1 + \frac{2(1-\sigma)}{\pi^2} \left(\log \frac{1+\sqrt{z}}{1-\sqrt{z}} \right)^2, & \text{if } \nu = 1 \\ 1 + \frac{2(1-\sigma)}{1-\nu^2} \sinh^2 \left[\left(\frac{2}{\pi} \arccos \nu \right) \operatorname{arc} \tanh \sqrt{z} \right], & \text{if } 0 < \nu < 1 \\ 1 + \frac{2(1-\sigma)}{1-\nu^2} \sin \left(\frac{\pi}{2R(t)} \int_0^{\frac{u(z)}{t}} \frac{1}{\sqrt{1-x^2}\sqrt{1-(tx)^2}} dx \right) + \frac{1}{\nu^2-1}, & \text{if } \nu > 1, \end{cases} \quad (5)$$

where $u(z) = \frac{z-\sqrt{t}}{1-\sqrt{tz}}$, $t \in (0, 1)$ and t is chosen such that $\nu = \cosh\left(\frac{\pi R'(t)}{4R(t)}\right)$, $R(t)$ is Legendre’s complete elliptic integral of the first kind and $R'(t)$ is the complementary integral of $R(t)$. Clearly, $\hat{p}_{\nu,\sigma}(z)$ is in \mathcal{P} with the expansion of the form

$$\hat{p}_{\nu,\sigma}(z) = 1 + \tau_1 z + \tau_2 z^2 + \dots, \quad (\tau_j = p_j(\nu, \sigma), j = 1, 2, 3, \dots), \quad (6)$$

we obtain

$$\tau_1 = \begin{cases} \frac{8(1-\sigma)(\arccos \nu)^2}{\pi^2(1-\nu^2)}, & \text{if } 0 \leq \nu < 1, \\ \frac{8(1-\sigma)}{\pi^2}, & \text{if } \nu = 1 \\ \frac{\pi^2(1-\sigma)}{4\sqrt{t}(v^2-1)R^2(t)(1+t)}, & \text{if } \nu > 1. \end{cases} \quad (7)$$

To avoid repetition, we let once for all throughout this paper

$$-1 \leq G < F \leq 1, 0 \leq \alpha < 1, -\frac{\pi}{2} < \theta < \frac{\pi}{2}, p, n \in \mathbb{N}, m \in \mathbb{N}_0.$$

Additionally, let

$$\chi(z) = d_0 + d_1 z + d_2 z^2 + \dots \quad (d_0 \neq 0) \quad \text{and} \quad |d_0| \leq 1. \quad (8)$$

Motivated by the study of Tang, Karthikeyan and Murugusundaramoorthy [23] and definition of $\mathcal{N}_p^n(\lambda, \beta, m; F, G)$, we now introduce the following class of functions:

Definition 1. For $t \in \mathbb{C}$, with $|t| \leq 1, t \neq 1, \lambda \geq 0$, and $\chi(z)$ is defined as in (8), we say that the function $h \in \mathcal{A}(p, 1)$ belongs to the class $\mathfrak{S}_p^m(b; \psi; \alpha; \lambda; F; G; \theta)$ if it satisfies the subordination condition

$$\frac{1 + i \tan \theta}{b} \left[\frac{Y_\lambda^p(m; t) z^{1-\lambda(p-m)} h^{(m+1)}(z)}{[h^{(m)}(z) - h^{(m)}(tz)]^{1-\lambda}} - p + m \right] \prec_{\kappa} \aleph(p; F, G; \alpha; \psi; z) - p \quad (9)$$

where p is an odd integer, $Y_\lambda^p(m; t) = (1 - t^p)^{(1-\lambda)} [\delta(p, m)]^{-\lambda}$ and $\aleph(p; F, G; \alpha; \psi; z)$ defined as in (4).

Remark 1. Now we will present some special cases of our class.

- (i) Let $p = b = 1, \alpha = m = \theta = \lambda = 0, \chi(z) = 1$ and $\psi = \hat{p}_{\nu,\sigma}(z)$ (see (5)) in Definition 1, then the class $\mathfrak{S}_p^m(b; \psi; \alpha; \lambda; F; G; \theta)$ reduces to class $k - \mathcal{US}(F, G, \sigma, t)$ defined by Arif et al. [24] (Definition 1.3) (also see [25]).
- (ii) If we replace $p = b = 1, \alpha = t = \theta = \lambda = 0, \chi(z) = 1$ and $\psi(z) = \hat{p}_{\nu,0}(z)$ in $\mathfrak{S}_p^m(b; \psi; \alpha; \lambda; F; G; \theta)$, where $\hat{p}_{\nu,0}(z)$ is defined as in (5), we can obtain $\eta - \mathcal{ST}[F, G]$ and $\eta - \mathcal{UC}[F, G]$ classes defined by Noor and Malik in [19] (Definition 1.3 and Definition 1.4) by choosing $m = 0$ and $m = 1$, respectively.
- (iii) If we let $\alpha = \lambda = m = \theta = 0, b = 1, p = 1, F = 1$ and $G = -1$, then $\mathfrak{S}_p^m(b; \psi; \alpha; \lambda; F; G; \theta)$ reduces to the classes $\mathcal{S}_*^s(\psi)$ defined by Shanmugam, Ramachandran and Ravichandran [26] (Definition 1.3).

(iv) If we let $t = \theta = m = \alpha = 0, p = b = 1, F = 1, G = -1$ and $\psi(z) = 1 + z/1 - z$, then the class $\mathfrak{S}_p^m(b; \psi; \alpha; \lambda; F; G; \theta)$ reduces to well-known class Bazilevič function defined by

$$\mathcal{B}(\lambda) = \left\{ h \in \mathcal{A}(1, 1); \operatorname{Re} \frac{z^{1-\lambda} h'(z)}{[h(z)]^{1-\lambda}} > 0 \right\}.$$

Apart from the above classes of functions, several classes of functions which were defined in earlier works are closer to the class of functions defined in Definition 1, for example see [21,27–33].

2. Preliminaries

In this section, we will state some results, which we will be using to establish our main results namely subordination properties and coefficient inequalities.

Lemma 1. Ref. [34] If $\vartheta(z) = 1 + \sum_{k=1}^{\infty} \vartheta_k z^k \in \mathcal{P}$, then $|\vartheta_k| \leq 2$ for all $k \geq 1$, and the inequality is sharp for $\vartheta_{\mu}(z) = \frac{1 + \mu z}{1 - \mu z}, |\mu| \leq 1$.

Lemma 2. Ref. [5] Let $\vartheta(z) = 1 + \sum_{k=1}^{\infty} \vartheta_k z^k \in \mathcal{P}$ and also let v be a complex number, then

$$|\vartheta_2 - v\vartheta_1^2| \leq 2 \max\{1, |2v - 1|\},$$

the result is sharp for functions given by

$$\vartheta(z) = \frac{1 + z^2}{1 - z^2}, \quad \vartheta(z) = \frac{1 + z}{1 - z}.$$

Lemma 3. Ref. [35] Let r be convex in \mathbb{E} , with $r(0) = a, \delta \neq 0$ and $\operatorname{Re} \delta \geq 0$. If $k \in \mathcal{H}(a, n)$ and

$$k(z) + \frac{zk'(z)}{\delta} \prec r(z),$$

then

$$k(z) \prec q(z) \prec r(z),$$

where

$$q(z) = \frac{\delta}{nz^{\delta/n}} \int_0^z r(t) t^{(\delta/n)-1} dt.$$

The function q is convex and is the best (a, n) -dominant.

Throughout this paper, we let

$$\aleph(p; F, G; \alpha; \psi; z) = \frac{[(1 + F)p + \alpha(G - F)]\psi(z) + [(1 - F)p - \alpha(G - F)]}{[(G + 1)\psi(z) + (1 - G)]} \tag{10}$$

From [18] (Theorem 2), with

$$w(z) = \frac{1}{2}\vartheta_1 z + \frac{1}{2}\left(\vartheta_2 - \frac{1}{2}\vartheta_1^2\right)z^2 + \frac{1}{2}\left(\vartheta_3 - \vartheta_1\vartheta_2 + \frac{1}{4}\vartheta_1^3\right)z^3 + \dots, z \in \mathbb{E},$$

we can obtain

$$\frac{b\chi(z)}{1+i\tan\theta} \{ \aleph(p; F, G; \alpha; \psi; w(z)) - p \} = \frac{bd_0L_1\vartheta_1(F-G)(p-\alpha)}{4(1+i\tan\theta)} z + \frac{b(F-G)(p-\alpha)d_0L_1}{4(1+i\tan\theta)} \left[\vartheta_2 - \vartheta_1^2 \left(\frac{(G+1)L_1+2\left(1-\frac{L_2}{L_1}\right)}{4} \right) + \frac{d_1\vartheta_1}{d_0} \right] z^2 + \dots \tag{11}$$

3. Fekete-Szegő Inequalities for the Class $\mathfrak{S}_p^m(b; \psi; \alpha; \lambda; F; G; \theta)$

Obtaining the solution to the Fekete-Szegő problem has been a main focus of researchers in this field, as it plays an very important role in obtaining the algebraic properties of a function. It continues to remain in spotlight to date, refer [36–38] where authors have obtained the Fekete-Szegő inequality for classes of functions with respect to symmetric points.

In this section, we obtain the solution to the Fekete-Szegő problem for functions belonging to the class $\mathfrak{S}_p^m(b; \psi; \alpha; \lambda; F; G; \theta)$.

Theorem 1. If $h(z) = z^p + \sum_{j=p+1}^{\infty} a_j z^j$, $p \in N = \{1, 2, 3, \dots\}$ and $h(z) \in \mathfrak{S}_p^m(b; \psi; \alpha; \lambda; F; G; \theta)$, then for odd values of $p + m$ we have

$$|a_{p+1}| \leq \frac{L_1 |b|(F - G)(p - \alpha) |\Gamma_1|}{2(p + 1) \sec \theta}, \tag{12}$$

and

$$|a_{p+2}| \leq \frac{L_1 |b|(F - G)(p - \alpha) |\Gamma_2|}{2 \sec \theta} \left[\left| \frac{d_1}{d_0} \right| + \max \left\{ 1, \left| \frac{(G+1)L_1}{2} - \frac{L_2}{L_1} - \frac{bd_0(F - G)(p - \alpha)L_1 \Gamma_1^2 \Gamma_3}{4(1 + i \tan \theta)} \right| \right\} \right], \tag{13}$$

where Γ_1, Γ_2 and Γ_3 are given by

$$\begin{aligned} \Gamma_1 &= \frac{(p - m + 1)(1 - t^p)}{(p - m + 1)(1 - t^p) + (p - m)(\lambda - 1)(1 - t^{p+1})} \\ \Gamma_2 &= \frac{(p - m + 1)(p - m + 2)(1 - t^p)}{(p + 1)(p + 2)[(p - m + 2)(1 - t^p) + (p - m)(\lambda - 1)(1 - t^{p+2})]} \\ \Gamma_3 &= \frac{(1 - t^{p+1})[2(p - m + 1)(1 - t^p)(\lambda - 1) + (p - m)(\lambda - 1)(\lambda - 2)(1 - t^{p+1})]}{(p - m + 1)^2(1 - t^p)^2}. \end{aligned}$$

In addition, for all $\mu \in \mathbb{C}$ we have

$$|a_{p+2} - \mu a_{p+1}^2| \leq \frac{L_1 |b|(F - G)(p - \alpha) |\Gamma_2|}{2 \sec \theta} \left[\left| \frac{d_1}{d_0} \right| + \max \{1, |2\mathcal{H}_1 - 1|\} \right],$$

where \mathcal{H}_1 is given by

$$\mathcal{H}_1 = \frac{1}{4} \left\{ (G + 1)L_1 + 2 \left(1 - \frac{L_2}{L_1} \right) - \frac{bd_0(F - G)(p - \alpha)L_1 \Gamma_1^2 \Gamma_3}{2(1 + i \tan \theta)} + \frac{\mu bd_0(F - G)(p - \alpha)L_1 \Gamma_1^2}{(p + 1)^2(1 + i \tan \theta)\Gamma_2} \right\}.$$

The inequality is sharp for each $\mu \in \mathbb{C}$.

Proof. By Definition 1, $h(z) \in \mathfrak{S}_p^m(b; \psi; \alpha; \lambda; F; G; \theta)$ implies

$$\frac{1 + i \tan \theta}{b} \left[\frac{\Gamma_\lambda^p(m; t) z^{1-\lambda(p-m)} h^{(m+1)}(z)}{[h^{(m)}(z) - h^{(m)}(tz)]^{1-\lambda}} - p + m \right] = \chi(z) [\aleph(p; F, G; \alpha; \psi; w(z)) - p], \tag{14}$$

where $\aleph(p; F, G; \alpha; \psi; w(z))$ is defined as in (10). For odd values of p , the left hand side of (14) is given by

$$\begin{aligned}
 & \frac{\Gamma_\lambda^p(m; t) z^{1-\lambda(p-m)} h^{(m+1)}(z)}{[h^{(m)}(z) - h^{(m)}(tz)]^{1-\lambda}} \\
 &= (p-m) \left\{ 1 + \left[\frac{(\lambda-1)(p+1)}{1!(p-m+1)} \left(\frac{1-t^{p+1}}{1-t^p} \right) + \frac{p+1}{p-m} \right] a_{p+1} z \right. \\
 &+ \left[\left(\frac{(\lambda-1)(p+1)(p+2)}{1!(p-m+1)(p-m+2)} \left(\frac{1-t^{p+2}}{1-t^p} \right) + \frac{(p+1)(p+2)}{(p-m+1)(p-m)} \right) a_{p+2} \right. \\
 &\quad \left. + \left(\frac{(\lambda-1)(\lambda-2)(p+1)^2}{2!(p-m+1)^2} \left(\frac{1-t^{p+1}}{1-t^p} \right)^2 + \right. \right. \\
 &\quad \left. \left. \frac{(\lambda-1)(p+1)^2}{1!(p-m)(p-m+1)} \left(\frac{1-t^{p+1}}{1-t^p} \right) \right) a_{p+1}^2 \right] z^2 + \dots \right\}. \tag{15}
 \end{aligned}$$

From (15) and (11), the coefficients of z and z^2 are given by

$$a_{p+1} = \frac{bd_0(F-G)(p-\alpha)L_1\vartheta_1\Gamma_1}{4(p+1)(1+i \tan \theta)} \tag{16}$$

and

$$a_{p+2} = \frac{bd_0(F-G)(p-\alpha)L_1\Gamma_2}{4(1+i \tan \theta)} \left[\vartheta_2 - \frac{\vartheta_1^2}{4} \left(L_1(G+1) + 2 \left(1 - \frac{L_2}{L_1} \right) - \frac{bd_0(F-G)(p-\alpha)L_1\Gamma_1^2\Gamma_3}{2(1+i \tan \theta)} \right) + \frac{d_1\vartheta_1}{d_0} \right]. \tag{17}$$

Applying Lemma 1 on (16), we can obtain (12). Using (17) together with Lemma 1, we have

$$\begin{aligned}
 |a_{p+2}| &= \frac{|b| |d_0|(F-G)(p-\alpha)L_1|\Gamma_2|}{4 \sec \theta} \left| \vartheta_2 - \frac{\vartheta_1^2}{4} \left((G+1)L_1 + 2 \left(1 - \frac{L_2}{L_1} \right) \right. \right. \\
 &\quad \left. \left. - \frac{bd_0(F-G)(p-\alpha)L_1\Gamma_1^2\Gamma_3}{2(1+i \tan \theta)} \right) + \frac{d_1\vartheta_1}{d_0} \right| \\
 &\leq \frac{|b|(F-G)(p-\alpha)L_1|\Gamma_2|}{2 \sec \theta} \left[\left| \frac{d_1}{d_0} \right| + \max \left\{ 1; \left| \frac{(G+1)L_1}{2} - \frac{L_2}{L_1} \right. \right. \right. \\
 &\quad \left. \left. \left. - \frac{bd_0(F-G)(p-\alpha)L_1\Gamma_1^2\Gamma_3}{4(1+i \tan \theta)} \right| \right\} \right].
 \end{aligned}$$

Hence the proof of (13).

Now to prove the Fekete-Szegő inequality for the class $\mathfrak{S}_p^m(b; \psi; \alpha; \lambda; F; G; \theta)$, we consider

$$\begin{aligned}
 |a_{p+2} - \mu a_{p+1}^2| &= \left| \frac{bd_0(F-G)(p-\alpha)L_1\Gamma_2}{4(1+i \tan \theta)} \left[\vartheta_2 - \frac{\vartheta_1^2}{4} \left((G+1)L_1 + 2 \left(1 - \frac{L_2}{L_1} \right) \right. \right. \right. \\
 &\quad \left. \left. \left. - \frac{bd_0(F-G)(p-\alpha)L_1\Gamma_1^2\Gamma_3}{2(1+i \tan \theta)} \right) + \frac{d_1\vartheta_1}{d_0} \right] - \frac{\mu d_0^2 b^2 (F-G)^2 (p-\alpha)^2 L_1^2 \vartheta_1^2 \Gamma_1^2}{16(p+1)^2(1+i \tan \theta)^2} \right| \\
 &= \left| \frac{bd_0(F-G)(p-\alpha)L_1\Gamma_2}{4(1+i \tan \theta)} \left[\vartheta_2 - \frac{\vartheta_1^2}{2} + \frac{\vartheta_1^2}{4} \left(\frac{2L_2}{L_1} - (G+1)L_1 \right. \right. \right. \\
 &\quad \left. \left. \left. + \frac{bd_0(F-G)(p-\alpha)L_1\Gamma_1^2\Gamma_3}{2(1+i \tan \theta)} - \frac{\mu b(F-G)(p-\alpha)L_1\Gamma_1^2}{(p+1)^2(1+i \tan \theta)\Gamma_2} \right) + \frac{d_1\vartheta_1}{d_0} \right] \right| \tag{18} \\
 &\leq \frac{|b|(F-G)(p-\alpha)|L_1|\Gamma_2}{4 \sec \theta} \left[2 + \frac{|\vartheta_1|^2}{4} \left(\left| \frac{2L_2}{L_1} - (G+1)L_1 \right. \right. \right. \\
 &\quad \left. \left. \left. + \frac{bd_0(F-G)(p-\alpha)L_1\Gamma_1^2\Gamma_3}{2(1+i \tan \theta)} - \frac{\mu bd_0(F-G)(p-\alpha)L_1\Gamma_1^2}{(p+1)^2(1+i \tan \theta)\Gamma_2} \right| - 2 \right) + 2 \left| \frac{d_1}{d_0} \right| \right].
 \end{aligned}$$

Denoting

$$H := \left| \frac{2L_2}{L_1} - (G+1)L_1 + \frac{bd_0(F-G)(p-\alpha)L_1\Gamma_1^2\Gamma_3}{2(1+i \tan \theta)} - \frac{\mu bd_0(F-G)(p-\alpha)L_1\Gamma_1^2}{(p+1)^2(1+i \tan \theta)\Gamma_2} \right|.$$

If $H \leq 2$, from (18) we obtain

$$|a_{p+2} - \mu a_{p+1}^2| \leq \frac{|b|(F-G)(p-\alpha)|L_1|\Gamma_2}{2 \sec \theta} \left| \frac{d_1}{d_0} \right|. \tag{19}$$

Further, if $H \geq 2$ from (18) we deduce

$$\begin{aligned}
 |a_{p+2} - \mu a_{p+1}^2| &\leq \frac{|b|(F-G)(p-\alpha)|L_1|\Gamma_2}{2 \sec \theta} \left(\left| \frac{2L_2}{L_1} - (G+1)L_1 \right. \right. \\
 &\quad \left. \left. + \frac{bd_0(F-G)(p-\alpha)L_1\Gamma_1^2\Gamma_3}{2(1+i \tan \theta)} - \frac{\mu bd_0(F-G)(p-\alpha)L_1\Gamma_1^2}{(p+1)^2(1+i \tan \theta)\Gamma_2} \right| + \left| \frac{d_1}{d_0} \right| \right). \tag{20}
 \end{aligned}$$

Equality of (19) will be attained if $\vartheta_1 = 0$, $\vartheta_2 = 2$ and $d_0 = 1$. Equivalently, by Lemma 2 we have $\psi(z^2) = \psi_2(z) = \frac{1+z^2}{1-z^2}$. Therefore, the extremal function of the class $\mathfrak{S}_p^m(b; \psi; \alpha; \lambda; F; G; \theta)$ is given by

$$\begin{aligned}
 &\frac{1+i \tan \theta}{b} \left[\frac{\Gamma_\lambda^p(m; t) z^{1-\lambda(p-m)} h^{(m+1)}(z)}{[h^{(m)}(z) - h^{(m)}(tz)]^{1-\lambda}} - p + m \right] \\
 &= \frac{[(1+F)p + \alpha(G-F)]\psi(z^2) + [(1-F)p - \alpha(G-F)]}{[(G+1)\psi(z^2) + (1-G)]} - p.
 \end{aligned}$$

Similarly, equality of (20) will be attained if $\vartheta_2 = 2$. Equivalently, by Lemma 2 we have $\psi(z) = \psi_1(z) = \frac{1+z}{1-z}$ and $\chi_1(z) = 1 + z + z^2 + \dots$. Therefore, the extremal function in $\mathfrak{S}_p^m(b; \psi; \alpha; \lambda; F; G; \theta)$ is given by

$$\begin{aligned}
 &1 + \frac{1+i \tan \alpha}{\gamma} \left[\frac{z^{1-t} [H'(z)]^\lambda}{[(1-\beta)H(z) + \beta z]^{1-t}} - 1 \right] \\
 &= \chi_1(z) \left[\frac{[(1+F)p + \alpha(G-F)]\psi_1(z) + [(1-F)p - \alpha(G-F)]}{[(G+1)\psi_1(z) + (1-G)]} - p \right],
 \end{aligned}$$

and the proof of the theorem is complete. \square

If we let $p = b = 1$, $\alpha = \theta = \lambda = 0$, $\chi(z) = 1$ and $\psi = \hat{p}_{v,\sigma}(z)$ and $m = 0$ in Theorem 1, we obtain the following result.

Corollary 1. Ref. [24] (Theorem 2.3) If $h(z) \in k - \mathcal{US}(F, G, \sigma, t)$ (see Remark 1 (i)), then we have

$$|a_2| \leq \frac{(F - G)|\tau_1|}{2|1 - t|},$$

and

$$|a_3| \leq \frac{(F - G)|\tau_1|}{2|2 - t - t^2|} \max \left\{ 1; \left| \frac{(G + 1)\tau_1}{2} - \frac{\tau_2}{\tau_1} + \frac{(F - G)\tau_1(1 + t)}{2(1 - t)} \right| \right\},$$

In addition, for all $\mu \in \mathbb{C}$ we have

$$|a_3 - \mu a_2^2| \leq \frac{(F - G)|\tau_1|}{2|1 - t - t^2|} \max \{1, |2\mathcal{H}_1 - 1|\}$$

where \mathcal{H}_1 is given by

$$\mathcal{H}_1 = \frac{1}{4} \left\{ (G + 1)\tau_1 + 2 \left(1 - \frac{\tau_2}{\tau_1} \right) + \frac{(F - G)\tau_1(1 + t)}{2(1 - t)} + \frac{\mu(F - G)\tau_1[2 - t - t^2]}{(1 - t)^2} \right\}.$$

The inequality is sharp for each $\mu \in \mathbb{C}$.

If we let $p = b = 1$, $\alpha = \theta = \lambda = 0$, $\chi(z) = 1$, $F = 1$, $G = -1$ and $m = 0$ in Theorem 1, we obtain the following result which was obtained by Shanmugam et al. [26] for real valued μ .

Corollary 2. Ref. [26] (Theorem 2.1)

If $h(z) \in \mathcal{S}_*^s(\psi)$ (see Remark 1 (iii)), then we have

$$|a_2| \leq L_1,$$

and

$$|a_3| \leq \frac{L_1}{2} \max \left\{ 1; \left| \frac{L_2}{L_1} - 4L_1 \right| \right\},$$

In addition, for all $\mu \in \mathbb{C}$ we have

$$|a_3 - \mu a_2^2| \leq \frac{L_1}{2} \max \left\{ 1, \left| \frac{L_2}{L_1} - 2L_1(1 + \mu) \right| \right\}.$$

The inequality is sharp for each $\mu \in \mathbb{C}$.

Some Applications Involving Bernoulli Lemniscate and Shell Shaped Region

Raina and Sokół [7] (also see [39]) defined the class $\mathcal{S}^*(z + \sqrt{1 + z^2})$. The function $\psi(z) = z + \sqrt{1 + z^2}$ maps the unit disc onto the interior of lune-shaped (shell-shaped) starlike region (see Figure 3a). The impact of $\aleph(p; F, G; \alpha; \psi; z)$ on the shell-shaped region is illustrated in Figure 3. It could be seen that if the distance between F and G are increased, then the mapping of unit disc becomes convex. If they are closer to each other, then the mapping is starlike. Furthermore, notice that in Figure 3, we have shown by the varying parameters involved that a shell-shaped region with corner $-2i$ is rotated to π radians in a counterclockwise direction and corner $+2i$ is rotated to π radians in clockwise direction.




Figure 3. Impact of $\aleph(p; F, G; \alpha; \psi; z)$ on the conic region $\psi(z) = z + \sqrt{1+z^2}$. (a) Mapping of \mathbb{E} under the transformation $\psi(z) = z + \sqrt{1+z^2}$. (b) Mapping of \mathbb{E} under the transformation $\aleph(1; 0, -0.3; 0.9; \psi; z)$ if $\psi(z) = z + \sqrt{1+z^2}$. (c) Mapping of \mathbb{E} under the transformation $\aleph(1; 0.9, 0.0; 0.5; \psi; z)$ if $\psi(z) = z + \sqrt{1+z^2}$. (d) Mapping of \mathbb{E} under the transformation $\aleph(1; 0.9, 0.8; 0.8; \psi; z)$ if $\psi(z) = z + \sqrt{1+z^2}$.

Corollary 3. Ref. [32] If $h \in S^*(z + \sqrt{1+z^2})$, then $|a_2| \leq 1$, $|a_3| \leq \frac{3}{4}$ and $|a_3 - \mu a_2^2| \leq \max\left\{\frac{1}{2}, \left|\mu - \frac{3}{4}\right|\right\}$.

Proof. The function $\psi(z) = z + \sqrt{1+z^2}$ has a Maclaurin series expansion of the form

$$\psi(z) = 1 + z + \frac{z^2}{2} - \frac{z^4}{8} + \frac{z^6}{16} - \frac{5z^8}{128} + \frac{7z^{10}}{256} - \frac{21z^{12}}{1024} + O[z]^{13}.$$

Now if we let $t = \alpha = \theta = \lambda = 0$, $p = F = 1$, $G = -1$, $L_1 = 1$ and $L_2 = \frac{1}{2}$ in Theorem 1, we obtain the assertion of the Corollary. \square

The function $\psi(z) = \sqrt{1+z}$ maps \mathbb{E} onto a set bounded by Bernoulli lemniscate (see [6]). Subfigures in Figure 4 describes the impact of $\aleph(p; F, G; \alpha; \psi; z)$ on Bernoulli lemniscate.

Corollary 4. Ref. [6] (Theorem 2) If $h \in S^*(\sqrt{1+z})$, then $|a_2| \leq \frac{1}{2}$, $|a_3| \leq \frac{1}{4}$ and $|a_3 - \mu a_2^2| \leq \max\left\{\frac{1}{4}, \left|\mu - \frac{7}{4}\right|\right\}$.

Proof. The function $\psi(z) = \sqrt{1+z}$ has a Maclaurin series expansion of the form

$$\psi(z) = 1 + \frac{z}{2} - \frac{z^2}{8} + \frac{z^3}{16} - \frac{5z^4}{128} + \frac{7z^5}{256} - \frac{21z^6}{1024} + \frac{33z^7}{2048} - \frac{429z^8}{32768} + O[z]^9.$$

Now if we let $t = \alpha = \theta = \lambda = 0$, $p = F = 1$, $G = -1$, $L_1 = \frac{1}{2}$ and $L_2 = -\frac{1}{8}$ in Theorem 1, we obtain the assertion of the Corollary. \square

Remark 2. By specializing the parameters involved, we can easily obtain the coefficient inequalities of starlike functions with respect to symmetric points associated with Bernoulli lemniscate and Shell-shaped region.




Figure 4. Impact of $\aleph(p; F, G; \alpha; \psi; z)$ on the conic region $\psi(z) = \sqrt{1+z}$. (a) Mapping of \mathbb{E} under the transformation $\psi(z) = \sqrt{1+z}$. (b) Mapping of \mathbb{E} under the transformation $\aleph(1; 0, -0.3; 0.9; \psi; z)$ if $\psi(z) = \sqrt{1+z}$. (c) Mapping of \mathbb{E} under the transformation $\aleph(1; 0.9, 0.0; 0.5; \psi; z)$ if $\psi(z) = \sqrt{1+z}$. (d) Mapping of \mathbb{E} under the transformation $\aleph(1; 0.9, 0.8; 0.8; \psi; z)$ if $\psi(z) = \sqrt{1+z}$.

4. Subordination Results for Functions with Respect to Symmetric Points

Researchers have investigated and obtained several interesting subordination conditions, see for example [20,21,40,41]. In this section we follow the steps detailed in Goyal

and Goswami [42], to obtain some sufficient conditions for functions to be in our defined function class. We let

$$\omega_\theta = (1 + i \tan \theta) \quad \text{and} \quad G^m(z, t) = [h^{(m)}(z) - h^{(m)}(tz)].$$

We begin with the following

Theorem 2. Let $h \in \mathcal{A}(p, 1)$ with $h^{(m)}(z)$, $h^{(m+1)}(z)$ and $G^m(z, t) \neq 0$ for all $z \in \mathbb{E} \setminus \{0\}$. Moreover, let $\aleph(p; F, G; \alpha; \psi; z)$ be convex univalent in \mathbb{E} with $\aleph(p; F, G; \alpha; 0) = p$ and $\text{Re } \aleph(p; F, G; \alpha; \psi; z) > 0$. Further suppose that

$$\left(p + \frac{\omega_\theta}{b\chi(z)} \left[\frac{Y_\lambda^p(m; t)z^{1-\lambda(p-m)}h^{(m+1)}(z)}{[G^m(z, t)]^{1-\lambda}} - p + m \right] \right)^2 \left[1 + 2 \left\{ \frac{(\lambda-1)zG^{m+1}(z, t)}{G^m(z, t)} \right. \right. \tag{21}$$

$$+ \frac{(1-\lambda)zG^{m+1}(z, t)[pb\chi(z) - (p-m)\omega_\theta][G^m(z, t)]^{-\lambda} + pbz\chi'(z)[G^m(z, t)]^{1-\lambda}}{[pb\chi(z) - (p-m)\omega_\theta][G^m(z, t)]^{1-\lambda} + \omega_\theta Y_\lambda^p(m; t)z^{1-\lambda(p-m)}h^{(m+1)}(z)}$$

$$+ \frac{\omega_\theta Y_\lambda^p(m; t)z^{1-\lambda(p-m)}[(1-\lambda(p-m))h^{(m+1)}(z) + zh^{(m+2)}(z)]}{[pb\chi(z) - (p-m)\omega_\theta][G^m(z, t)]^{1-\lambda} + \omega_\theta Y_\lambda^p(m; t)z^{1-\lambda(p-m)}h^{(m+1)}(z)}$$

$$\left. - \frac{z\chi'(z)}{\chi(z)} \right\} \prec \aleph(p; F, G; \alpha; \psi; z). \tag{22}$$

Then

$$\frac{1 + i \tan \theta}{b} \left[\frac{Y_\lambda^p(m; t)z^{1-\lambda(p-m)}h^{(m+1)}(z)}{[h^{(m)}(z) - h^{(m)}(tz)]^{1-\lambda}} - p + m \right] \prec_\kappa \phi(z) = \sqrt{Q(z)} - p \tag{23}$$

where

$$Q(z) = \frac{1}{z} \int_0^z \aleph(p; F, G; \alpha; \psi; t) dt$$

and ϕ is convex and is the best dominant.

Proof. Let

$$k(z) = p + \frac{1 + i \tan \theta}{b\chi(z)} \left[\frac{Y_\lambda^p(m; t)z^{1-\lambda(p-m)}h^{(m+1)}(z)}{[h^{(m)}(z) - h^{(m)}(tz)]^{1-\lambda}} - p + m \right] \quad (z \in \mathbb{E}),$$

then $k(z) \in \mathcal{H}(p, 1)$ with $k(z) \neq 0$.

Since $\aleph(p; F, G; \alpha; \psi; z)$ is convex, it can be easily seen that Q is convex and univalent in \mathbb{E} . If we make the change of the variables $K(z) = k^2(z)$, then $K(z) \in \mathcal{H}(p, 1)$ with $K(z) \neq 0$ in \mathbb{E} .

By a straight forward computation, we have

$$\frac{zK'(z)}{K(z)} = 2 \left[\frac{\omega_\theta Y_\lambda^p(m; t)z^{1-\lambda(p-m)}[(1-\lambda(p-m))h^{(m+1)}(z) + zh^{(m+1)}(z)]}{[pb\chi(z) - (p-m)\omega_\theta][G^m(z, t)]^{1-\lambda} + \omega_\theta Y_\lambda^p(m; t)z^{1-\lambda(p-m)}h^{(m+1)}(z)} \right.$$

$$+ \frac{(1-\lambda)zG^{m+1}(z, t)[pb\chi(z) - (p-m)\omega_\theta][G^m(z, t)]^{-\lambda} + pbz\chi'(z)[G^m(z, t)]^{1-\lambda}}{[pb\chi(z) - (p-m)\omega_\theta][G^m(z, t)]^{1-\lambda} + \omega_\theta Y_\lambda^p(m; t)z^{1-\lambda(p-m)}h^{(m+1)}(z)}$$

$$\left. - \frac{z\chi'(z)}{\chi(z)} + \frac{(\lambda-1)zG^{m+1}(z, t)}{G^m(z, t)} \right].$$

Thus, by (22), we have

$$K(z) + zK'(z) \prec r(z) \quad (z \in \mathbb{E}). \tag{24}$$

Now by Lemma 3, we deduce that

$$K(z) \prec Q(z) \prec r(z).$$

Since $Re r(z) > 0$ and $Q(z) \prec r(z)$ we also have $Re Q(z) > 0$. Hence the univalence of Q implies the univalence of $\sqrt{Q(z)}$ and $k^2(z) \prec Q(z)$ implies that $k(z) \prec \sqrt{Q(z)}$. Since subordination is invariant under translation and using the fact that $g/\chi \prec r$ implies $g \prec_{\kappa} r$, we have

$$\frac{1 + i \tan \theta}{b} \left[\frac{Y_{\lambda}^p(m; t) z^{1-\lambda(p-m)} h^{(m+1)}(z)}{[h^{(m)}(z) - h^{(m)}(tz)]^{1-\lambda}} - p + m \right] \prec_{\kappa} \sqrt{Q(z)} - p,$$

and the proof is complete. \square

If we let $p = b = 1, \lambda = m = \theta = 0$ in Theorem 2, we have

Corollary 5. Let $h \in \mathcal{A}$ with $h(z), h'(z)$ and $[h(z) - h(tz)] \neq 0$ for all $z \in \mathbb{E} \setminus \{0\}$. Furthermore, let $\aleph(p; F, G; \alpha; \psi; z)$ is convex in \mathbb{E} with $\aleph(p; F, G; \alpha; \psi; 0) = p$ and $Re \aleph(p; F, G; \alpha; \psi; z) > 0$. Further suppose that

$$\begin{aligned} & \left(1 + \frac{1}{\chi(z)} \left[\frac{zh'(z)}{[h(z) - h(tz)]} - 1 \right] \right)^2 \left[1 + 2 \left\{ \frac{zh'(z) + z^2 h''(z)}{[\chi(z) - 1][h(z) - h(tz)] + zh'(z)} \right. \right. \\ & \quad + \frac{z[h(z) - h(tz)]' [\chi(z) - 1] + z\chi'(z)[h(z) - h(tz)]}{[\chi(z) - 1][h(z) - h(tz)] + zh'(z)} \\ & \quad \left. \left. - \frac{z[h(z) - h(tz)]'}{[h(z) - h(tz)]} - \frac{z\chi'(z)}{\chi(z)} \right\} \right] \prec \aleph(p; F, G; \alpha; \psi; z). \end{aligned}$$

Then

$$\frac{zh'(z)}{[h(z) - h(tz)]} - 1 \prec_{\kappa} \phi(z) = \sqrt{Q(z)} - 1$$

where

$$Q(z) = \frac{1}{z} \int_0^z \aleph(p; F, G; \alpha; \psi; t) dt$$

and ϕ is convex and is the best dominant.

From the Corollary 5, we deduce that on letting $p = b = 1, m = \alpha = \theta = \lambda = 0, \chi(z) = 1$ and $\psi = \hat{p}_{\nu, \sigma}(z)$ (see (5)) in Theorem 1, then we can obtain the sufficient conditions for functions to be in $k - \mathcal{US}(F, G, \sigma, t)$ (see Remark 1 (i)).

Corollary 6. Let $h \in \mathcal{A}$ with $h'(z)$ and $[h(z) - h(-z)] \neq 0$ for all $z \in \mathbb{E} \setminus \{0\}$. If

$$Re \left\{ \left(\frac{zh'(z)}{[h(z) - h(-z)]} \right)^2 \left[3 + \frac{2zh''(z)}{h'(z)} - \frac{2z[h(z) - h(-z)]'}{[h(z) - h(-z)]} \right] \right\} > \alpha,$$

then

$$Re \frac{zh'(z)}{[h(z) - h(-z)]} > \eta(\alpha),$$

where $\eta(\alpha) = [2(1 - \alpha) \cdot \log 2 + (2\alpha - 1)]^{\frac{1}{2}}$. This result is sharp

Proof. Letting $p = 1, F = 1, G = -1$ and $\psi(z) = \frac{1-z}{1+z}$ in (4), we obtain

$$\aleph(1; 1, -1; \alpha; z) = \frac{1 + (2\alpha - 1)z}{1 + z}, \quad (0 \leq \alpha < 1).$$

Additionally, if we let $\chi(z) = 1, p = b = 1, \theta = m = \lambda = 0$, in Theorem 2, we have

$$Q(z) = \frac{1}{z} \int_0^z \frac{1 + (2\alpha - 1)t}{1 + t} dt$$

which is convex in \mathbb{E} along with $\operatorname{Re} Q(z) > 0$. Therefore

$$\min_{|z| \leq 1} \operatorname{Re} \sqrt{Q(z)} = \sqrt{Q(1)} = [2(1 - \alpha) \cdot \log 2 + (2\alpha - 1)]^{\frac{1}{2}}.$$

Hence the proof of the Corollary. \square

5. Classes of Multivalent Functions Using Quantum Calculus

Now, we give a very brief introduction of the q -calculus. We let

$$[n]_q = \sum_{k=1}^n q^{k-1}, \quad [0]_q = 0, \quad (q \in \mathbb{C}).$$

Srivastava in [43] initiated the study of geometric function theory in dual with quantum calculus in 1988. However, this duality theory was brought into the spotlight by Ismail et al. [44] who introduced and studied the so-called class of q -starlike functions. For detailed study of the developments and applications of this duality theory, refer to the recent survey-cum-expository article of Srivastava [45] and references provided therein.

The q -difference operator for a function $h \in \mathcal{A}(p, 1)$ is defined by

$$\mathfrak{D}_q h(z) := \begin{cases} h'(0), & \text{if } z = 0, \\ \frac{h(z) - h(qz)}{(1 - q)z}, & \text{if } z \neq 0. \end{cases} \tag{25}$$

From (25), if $h \in \mathcal{A}(p, 1)$ we can easily see that $\mathfrak{D}_q h(z) = pz^{p-1} + \sum_{k=p+n}^{\infty} [k]_q a_k z^{k-1}$, for $z \neq 0$ and note that $\lim_{q \rightarrow 1^-} \mathfrak{D}_q h(z) = h'(z)$. The q -Jackson integral is defined by (see [46])

$$I_q[h(z)] := \int_0^z h(t) d_q t = z(1 - q) \sum_{n=0}^{\infty} q^n h(zq^n) \tag{26}$$

provided the q -series converges. Further observe that

$$\mathfrak{D}_q I_q h(z) = h(z) \quad \text{and} \quad I_q \mathfrak{D}_q h(z) = h(z) - h(0),$$

where the second equality holds if h is continuous at $z = 0$. Ismail et al. in [44] defined the class \mathcal{S}_q^* as class of functions which satisfies the condition

$$\left| \frac{z \mathfrak{D}_q h(z)}{h(z)} - \frac{1}{1 - q} \right| \leq \frac{1}{1 - q}, \quad (h \in \mathcal{S}).$$

The class \mathcal{S}_q^* is the so-called class of q -starlike functions. Equivalently, a function $h \in \mathcal{S}_q^*$, if and only if the subordination condition (see ([47], Definition 7))

$$\frac{z \mathfrak{D}_q h(z)}{h(z)} \prec \frac{1 + z}{1 - qz},$$

holds.

Let us define the q -analogue of $\aleph(p; F, G; \alpha; \psi; z)$ (see (4)) as

$$\aleph_q(p; F, G; \alpha; \psi; z) = \frac{[(F + 1)[p]_q + \alpha(G - F)]\psi(z) + [(1 - F)[p]_q - \alpha(G - F)]}{[(G + 1)\psi(z) + (1 - G)]} \tag{27}$$

Srivastava et al. [47–54] introduced function classes of q -starlike functions related with conic region and also studied the impact of Janowski functions on those conic regions. For recent advances pertaining to quantum calculus, refer to Aldawish and Ibrahim [55] and Zhou et al. [56]. Motivated by aforementioned works on q -calculus, we define the following class by replacing ordinary derivative with q -derivative in function class $\mathfrak{S}_p^m(b; \psi; \alpha; \lambda; F; G; \theta)$ (see Definition 1).

Definition 2. Let $\mathfrak{D}_q^m h = \mathfrak{D}_q^{m-1}(\mathfrak{D}_q^1 h(z))$. For $t \in \mathbb{C}$, with $|t| \leq 1$, $\lambda \geq 0$, $p, n \in \mathbb{N}$, $m \in \mathbb{N}_0$ and $\aleph_q(p; F, G; \alpha; \psi; z)$ defined as in (27), we say that the function $h \in \mathcal{A}(p, 1)$ belongs to the class $\mathcal{QS}_p^m(b; \psi; \alpha; \lambda; \delta; F; G; \theta)$ if it satisfies the subordination condition

$$\frac{1 + i \tan \theta}{b} \left(\frac{\Gamma_\lambda^p(m; t) z^{1-\lambda(p-m)} \mathfrak{D}_q^{m+1} h(z)}{[\mathfrak{D}_q^m h(z) - \mathfrak{D}_q^m h(tz)]^{1-\lambda}} - [p - m]_q \right) \prec_\kappa \aleph_q(p; F, G; \alpha; \psi; z) - [p]_q \tag{28}$$

where $\Gamma_\lambda^p(m; t) = (1 - t^p)^{(1-\lambda)} [\Delta_q(p, m - 1)]^{-\lambda}$, $\psi \in \mathcal{P}$ is defined as in (2).

Suppose \prec_κ is replaced with \prec and let $\psi = \frac{1+z}{1-qz}$, $q \in (0, 1)$ in $\mathcal{QS}_p^m(b; \psi; \alpha; \lambda; \delta; F; G; \theta)$, then by definition of subordination of analytic function, a function $h \in \mathcal{A}(p, 1)$ is said to be in $\mathcal{QS}_p^m(b; \psi; \alpha; \lambda; \delta; F; G; \theta)$ if and only if ($q \in (0, 1)$, $z \in \mathbb{E}$),

$$\begin{aligned} & [p]_q + \frac{1 + i \tan \theta}{b} \left(\frac{\Gamma_\lambda^p(m; t) z^{1-\lambda(p-m)} \mathfrak{D}_q^{m+1} h(z)}{[\mathfrak{D}_q^m h(z) - \mathfrak{D}_q^m h(tz)]^{1-\lambda}} - [p - m]_q \right) \\ &= \frac{\{(F + 1)[p]_q + \alpha(G - F)\}w(z) + 2[p]_q + \{\alpha(G - F) - (1 - F)[p]_q\}qw(z)}{(G + 1)w(z) + 2 + (G - 1)qw(z)}, \end{aligned}$$

where $w(z)$ is analytic in \mathbb{E} and $w(0) = 0, |w(z)| < 1$.

Remark 3. If we let $m = \alpha = \lambda = \theta = 0$, $b = 1$, $p = 1$, $F = 1$ and $G = -1$, then $\mathcal{QS}_p^m(b; \psi; \alpha; \lambda; \delta; F; G; \theta)$ reduces to the classes $\mathcal{S}_{q,s}^*(\psi)$ defined by Ramachandran et al. [57] (Definition 1).

Main Results Involving Quantum calculus

We just state q -analogue result of Theorems 1 and 2. Here we have omitted the proof, as it could be obtained by retracing the steps of Theorems 1 and 2.

Theorem 3. If $h(z) = z^p + \sum_{j=p+1}^\infty a_j z^j$, $p \in \mathbb{N} = \{1, 2, 3, \dots\}$ and $h(z) \in \mathcal{QS}_p^m(b; \psi; \alpha; \lambda; \delta; F; G; \theta)$, then for odd values of p we have

$$|a_{p+1}| \leq \frac{|b|(F - G)(p - \alpha)|L_1 Y_1|}{2[p + 1]_q \sec \theta}, \tag{29}$$

and

$$|a_{p+2}| \leq \frac{|b|(F - G)(p - \alpha)|L_1 Y_2|}{2 \sec \theta} \left[\left| \frac{d_1}{d_0} \right| + \max \left\{ 1; \left| \frac{(G + 1)L_1}{2} - \frac{L_2}{L_1} - \frac{bd_0(F - G)(p - \alpha)L_1 Y_1^2 Y_3}{4(1 + i \tan \theta)} \right| \right\} \right],$$

where Y_1, Y_2 and Y_3 are given by

$$Y_1 = \frac{[p - m + 1]_q(1 - t^p)}{[p - m + 1]_q(1 - t^p) + [p - m]_q(\lambda - 1)(1 - t^{p+1})}$$

$$Y_2 = \frac{[p - m + 1]_q[p - m + 2]_q(1 - t^p)}{[p + 1]_q[p + 2]_q[[p - m + 2]_q(1 - t^p) + [p - m]_q(\lambda - 1)(1 - t^{p+2})]}$$

$$Y_3 = \frac{(1 - t^{p+1})[2[p - m + 1]_q(1 - t^p)(\lambda - 1) + [p - m]_q(\lambda - 1)(\lambda - 2)(1 - t^{p+1})]}{[p - m + 1]_q^2(1 - t^p)^2}.$$

Furthermore, for all $\mu \in \mathbb{C}$ we have

$$|a_{p+2} - \mu a_{p+1}^2| \leq \frac{|b|(F - G)(p - \alpha)|L_1 Y_2|}{2 \sec \theta} \left[\left| \frac{d_1}{d_0} \right| + \max \{1, |2Q_1 - 1|\} \right],$$

where Q_1 is given by

$$Q_1 = \frac{1}{4} \left\{ (G + 1)L_1 + 2 \left(1 - \frac{L_2}{L_1} \right) - \frac{bd_0(F - G)(p - \alpha)L_1 Y_1^2 Y_3}{2(1 + i \tan \theta)} + \frac{\mu bd_0(F - G)(p - \alpha)L_1 Y_1^2}{(p + 1)^2(1 + i \tan \theta)Y_2} \right\}.$$

The inequality is sharp for each $\mu \in \mathbb{C}$.

Remark 4. If we let $q \rightarrow 1^-$ in Theorem 3, then we obtain the solution to the Fekete-Szegő problem of the class $\mathfrak{S}_p^m(b; \psi; \alpha; \lambda; F; G; \theta)$.

q -analogue of the Lemma 3 (ordinary derivative replaced with a quantum derivative) need not be true for all $q \in (0, 1)$. It is true only if we could choose a sequence q_n that tends to 1^- . Thus, we will use same lemma with ordinary derivative to establish the sufficient conditions for functions in $\mathcal{QS}_p^m(b; \psi; \alpha; \lambda; \delta; F; G; \theta)$.

Theorem 4. Let $h \in \mathcal{A}(p, 1)$ with $\mathfrak{D}_q^m h(z), \mathfrak{D}_q^{m+1} h(z)$ and $[\mathfrak{D}_q^m h(z) - \mathfrak{D}_q^m h(tz)] \neq 0$ for all $z \in \mathbb{E} \setminus \{0\}$. Furthermore, let $\aleph_q(p; F, G; \alpha; \psi; z)$ is convex in \mathbb{E} with $\aleph_q(p; F, G; \alpha; 0) = [p]_q$ and $\text{Re } \aleph_q(p; F, G; \alpha; \psi; z) > 0$. Let $\mathcal{G}_q^m(t; z) = \mathfrak{D}_q^m h(z) - \mathfrak{D}_q^m h(tz), \omega_\theta = 1 + i \tan \theta$ and $L(p; b; \chi; \theta) = ([p]_q b \chi(z) - [p - m]_q \omega_\theta)$. Further suppose that

$$\left[[p]_q + \frac{\omega_\theta}{b} \left(\frac{\Gamma_\lambda^p(m; t) z^{1-\lambda(p-m)} \mathfrak{D}_q^{m+1} h(z)}{[\mathcal{G}_q^m(t; z)]^{1-\lambda}} - [p-m]_q \right) \right]^2 \left[1 + 2 \left\{ \frac{(\lambda-1)z [\mathcal{G}_q^m(t; z)]'}{[\mathcal{G}_q^m(t; z)]} \right. \right. \\ \left. \left. + \frac{(1-\lambda)z [\mathcal{G}_q^m(t; z)]' L(p; b; \chi; \theta) [\mathcal{G}_q^m(t; z)]^{-\lambda} + [p]_q b z \chi'(z) [\mathcal{G}_q^m(t; z)]^{1-\lambda}}{L(p; b; \chi; \theta) [\mathcal{G}_q^m(t; z)]^{1-\lambda} + \omega_\theta \Gamma_\lambda^p(m; t) z^{1-\lambda(p-m)} (\mathfrak{D}_q^{m+1} h(z))'} \right. \right. \\ \left. \left. + \frac{\omega_\theta \Upsilon_\lambda^p(m; t) z^{1-\lambda(p-m)} ([1-\lambda(p-m)]_q \mathfrak{D}_q^{m+1} h(z) + z (\mathfrak{D}_q^{m+1} h(z))')}{L(p; b; \chi; \theta) [\mathcal{G}_q^m(t; z)]^{1-\lambda} + \omega_\theta \Gamma_\lambda^p(m; t) z^{1-\lambda(p-m)} (\mathfrak{D}_q^{m+1} h(z))'} \right. \right. \\ \left. \left. - \frac{z \chi'(z)}{\chi(z)} \right\} \right] \prec \aleph_q(p; F, G; \alpha; \psi; z).$$

Then

$$\frac{\omega_\theta}{b} \left(\frac{\Gamma_\lambda^p(m; t) z^{1-\lambda(p-m)} \mathfrak{D}_q^{m+1} h(z)}{[\mathcal{G}_q^m(t; z)]^{1-\lambda}} - [p-m]_q \right) \prec_\kappa \phi(z) = \sqrt{R(z)} - [p]_q$$

where

$$R(z) = \frac{1}{z} \int_0^z \aleph_q(p; F, G; \alpha; \psi; t) dt$$

and ϕ is convex and is the best dominant.

Remark 5. As $q \rightarrow 1^-$, the Theorem 4 reduces to Theorem 2.

6. Conclusions

The study of geometrical implications is an integral part of research in geometric function theory. Here we have shown that a function $\aleph(p; F, G; \alpha; z)$ which was defined analytically in [18] indeed has beautiful geometric implications.

Extension and unification of various well-known classes of functions were the main objective of this paper. We defined a new family of multivalent functions of complex order using higher order derivatives. Inclusion relations, Fekete-Szegő inequalities and subordination conditions for starlikeness of the defined function class have been established. Attempting discretization of the results, we extend the defined function class using q -derivative. All the results involving quantum calculus were just stated, as the method of proof though cumbersome but is similar to our main results.

Author Contributions: Conceptualization, K.R.K., S.L., S.V., E.U. and D.M.; methodology, K.R.K., S.L., S.V., E.U. and D.M.; software, K.R.K., S.V. and D.M.; validation, K.R.K., S.L., S.V., E.U. and D.M.; formal analysis, K.R.K., S.L., S.V., E.U. and D.M.; investigation, K.R.K., S.L., S.V., E.U. and D.M.; resources, K.R.K., S.L., S.V., E.U. and D.M.; data curation, K.R.K., S.L., S.V., E.U. and D.M.; writing—original draft preparation, S.V., E.U. and D.M.; writing—review and editing, K.R.K., S.L., S.V., E.U. and D.M.; visualization, K.R.K., S.L., S.V., E.U. and D.M.; supervision, K.R.K., S.V. and E.U.; project administration, K.R.K. All authors have read and agreed to the published version of the manuscript.

Funding: The APC was partially funded by National University of Science and Technology, Oman.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: Teodor Bulboacă provided detailed insights on the effects of replacing the quantum derivative with an ordinary derivative in Lemma 3, while finalizing the results of [21]. The first author would like to thank Teodor Bulboacă for providing his valuable comments, which led to the presumably correct version of Theorem 4 herein.

Conflicts of Interest: All the authors declare that they have no conflict of interest.

References

1. Goodman, A.W. *Univalent Functions*; Mariner Publishing Co., Inc.: Tampa, FL, USA, 1983; Volume I.
2. Hayman, W.K. *Multivalent Functions*; Cambridge Tracts in Mathematics and Mathematical Physics, No. 48; Cambridge University Press: Cambridge, UK, 1958.
3. Haji Mohd, M.; Darus, M. Fekete-Szegő problems for quasi-subordination classes. *Abstr. Appl. Anal.* **2012**, *2012*, 192956. [[CrossRef](#)]
4. Karthikeyan, K.R.; Murugusundaramoorthy, G.; Cho, N.E. Some inequalities on Bazilevič class of functions involving quasi-subordination. *AIMS Math.* **2021**, *6*, 7111–7124. [[CrossRef](#)]
5. Ma, W.C.; Minda, D. A unified treatment of some special classes of univalent functions. In *Lecture Notes Analysis, I, Proceedings of the Conference on Complex Analysis, Tianjin, China, 19–23 June 1992*; International Press Inc.: Cambridge, MA, USA, 1992; pp. 157–169.
6. Sokół, J. Coefficient estimates in a class of strongly starlike functions. *Kyungpook Math. J.* **2009**, *49*, 349–353. [[CrossRef](#)]
7. Raina, R.K.; Sokół, J. Some properties related to a certain class of starlike functions. *C. R. Math. Acad. Sci. Paris* **2015**, *353*, 973–978. [[CrossRef](#)]
8. Raina, R.K.; Sokół, J. Fekete-Szegő problem for some starlike functions related to shell-like curves. *Math. Slovaca* **2016**, *66*, 135–140. [[CrossRef](#)]
9. Aouf, M.K.; Dziok, J.; Sokół, J. On a Subclass of Strongly Starlike Functions. *Appl. Math. Lett.* **2011**, *24*, 27–32. [[CrossRef](#)]
10. Dziok, J.; Raina, R.K.; Sokół, J. On α -convex functions related to shell-like functions connected with Fibonacci numbers. *Appl. Math. Comput.* **2011**, *218*, 996–1002. [[CrossRef](#)]
11. Dziok, J.; Raina, R.K.; Sokół, J. Certain results for a class of convex functions related to a shell-like curve connected with Fibonacci numbers. *Comput. Math. Appl.* **2011**, *61*, 2605–2613. [[CrossRef](#)]
12. Dziok, J.; Raina, R.K.; Sokół, J. On a class of starlike functions related to a shell-like curve connected with Fibonacci numbers. *Math. Comput. Model.* **2013**, *57*, 1203–1211. [[CrossRef](#)]
13. Gandhi, S.; Ravichandran, V. Starlike functions associated with a lune. *Asian-Eur. J. Math.* **2017**, *10*, 1750064. [[CrossRef](#)]
14. Khatter, K.; Ravichandran, V.; Sivaprasad Kumar, S. Starlike functions associated with exponential function and the lemniscate of Bernoulli. *Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM* **2019**, *113*, 233–253. [[CrossRef](#)]
15. Mendiratta, R.; Nagpal, S.; Ravichandran, V. A subclass of starlike functions associated with left-half of the lemniscate of Bernoulli. *Internat. J. Math.* **2014**, *25*, 1450090. [[CrossRef](#)]
16. Janowski, W. Some extremal problems for certain families of analytic functions I. *Ann. Polon. Math.* **1973**, *10*, 297–326. [[CrossRef](#)]
17. Aouf, M.K. On a class of p -valent starlike functions of order α . *Internat. J. Math. Math. Sci.* **1987**, *10*, 733–744. [[CrossRef](#)]
18. Breaz, D.; Karthikeyan, K.R.; Senguttuvan, A. Multivalent prestarlike functions with respect to symmetric points. *Symmetry* **2022**, *14*, 20.
19. Noor, K.I.; Malik, S.N. On coefficient inequalities of functions associated with conic domains. *Comput. Math. Appl.* **2011**, *62*, 2209–2217. [[CrossRef](#)]
20. Aouf, M.K.; Bulboacă, T.; Seoudy, T.M. Subclasses of multivalent non-Bazilevič functions defined with higher order derivatives. *Bull. Transilv. Univ. Braşov Ser. III* **2020**, *13*, 411–422. [[CrossRef](#)]
21. Karthikeyan, K.R.; Murugusundaramoorthy, G.; Bulboacă, T. Properties of λ -pseudo-starlike functions of complex order defined by subordination. *Axioms* **2021**, *10*, 86. [[CrossRef](#)]
22. Ahuja, O.; Bohra, N.; Cetinkaya, A.; Kumar, S. Univalent functions associated with the symmetric points and cardioid-shaped domain involving (p, q) -calculus. *Kyungpook Math. J.* **2021**, *61*, 75–98.
23. Tang, H.; Karthikeyan, K.R.; Murugusundaramoorthy, G. Certain subclass of analytic functions with respect to symmetric points associated with conic region. *AIMS Math.* **2021**, *6*, 12863–12877. [[CrossRef](#)]
24. Arif, M.; Wang, Z.-G.; Khan, M.R.; Lee, S.K. Coefficient inequalities for Janowski-sakaguchi type functions associated with conic regions. *Hacet. J. Math. Stat.* **2018**, *47*, 261–271. [[CrossRef](#)]
25. Arif, M.; Ahmad, K.; Liu, J.-L.; Sokół, J. A new class of analytic functions associated with Sălăgean operator. *J. Funct. Spaces* **2019**, *8*, 6157394. [[CrossRef](#)]
26. Shanmugam, T.N.; Ramachandran, C.; Ravichandran, V. Fekete-Szegő problem for subclasses of starlike functions with respect to symmetric points. *Bull. Korean Math. Soc.* **2006**, *43*, 589–598. [[CrossRef](#)]
27. Ibrahim, R.W. On a Janowski formula based on a generalized differential operator. *Commun. Fac. Sci. Univ. Ank. Ser. A1 Math. Stat.* **2020**, *69*, 1320–1328.
28. Kavitha, D.; Dhanalakshmi, K. Subclasses of analytic functions with respect to symmetric and conjugate points bounded by conical domain. *Adv. Math. Sci. J.* **2020**, *9*, 397–404. [[CrossRef](#)]

29. Mohankumar, D.; Senguttuvan, A.; Karthikeyan, K.R.; Ganapathy Raman, R. Initial coefficient bounds and Fekete-Szegő problem of pseudo-Bazilevič functions involving quasi-subordination. *Adv. Dyn. Syst. Appl.* **2021**, *16*, 767–777.
30. Mashwan, W.K.; Ahmad, B.; Khan, M.G.; Mustafa, S.; Arjika, S.; Khan, B. Pascu-Type analytic functions by using Mittag-Leffler functions in Janowski domain. *Math. Probl. Eng.* **2021**, *2021*, 1209871. [[CrossRef](#)]
31. Raina, R.K.; Sokół, J. On a class of analytic functions governed by subordination. *Acta Univ. Sapientiae Math.* **2019**, *11*, 144–155. [[CrossRef](#)]
32. Raina, R.K.; Sokół, J. On coefficient estimates for a certain class of starlike functions. *Haceppt. J. Math. Stat.* **2015**, *44*, 1427–1433. [[CrossRef](#)]
33. Sokół, J.; Thomas, D.K. Further results on a class of starlike functions related to the Bernoulli lemniscate. *Houst. J. Math.* **2018**, *44*, 83–95.
34. Pommerenke, C. *Univalent Functions*; Vandenhoeck & Ruprecht: Göttingen, Germany, 1975; p. 376.
35. Hallenbeck, D.J.; Ruscheweyh, S. Subordination by convex functions. *Proc. Amer. Math. Soc.* **1975**, *52*, 191–195. [[CrossRef](#)]
36. Breaz, D.; Cotîrlă, L.-I. The study of the new classes of m -Fold symmetric bi-univalent functions. *Mathematics* **2022**, *10*, 75. [[CrossRef](#)]
37. Oros, G.I.; Cotîrlă, L.-I. Coefficient estimates and the Fekete–Szegő problem for new classes of m -fold symmetric bi-univalent functions. *Mathematics* **2022**, *10*, 129. [[CrossRef](#)]
38. Srivastava, H.M.; Kamali, M.; Urdaletova, A. A study of the Fekete-Szegő functional and coefficient estimates for subclasses of analytic functions satisfying a certain subordination condition and associated with the Gegenbauer polynomials. *AIMS Math.* **2022**, *7*, 2568–2584. [[CrossRef](#)]
39. Murugusundaramoorthy, G.; Bulboacă, T. Hankel determinants for new subclasses of analytic functions related to a shell shaped region. *Mathematics* **2020**, *8*, 1041. [[CrossRef](#)]
40. Ibrahim, R.W.; Baleanu, D. Analytic solution of the Langevin differential equations dominated by a multibrot fractal set. *Fractal Fract.* **2021**, *5*, 50. [[CrossRef](#)]
41. Ibrahim, R.W.; Baleanu, D. On quantum hybrid fractional conformable differential and integral operators in a complex domain. *Rev. Real Acad. Cienc. Exactas Fís. Natur. Ser. A Mat. (RACSAM)* **2021**, *31*, 115. [[CrossRef](#)]
42. Goyal, S.P.; Goswami, P. On sufficient conditions for analytic functions to be Bazilevič. *Complex Var. Elliptic Equ.* **2009**, *54*, 485–492. [[CrossRef](#)]
43. Srivastava, H.M. Univalent functions, fractional calculus, and associated generalized hypergeometric functions. In *Univalent Functions, Fractional Calculus, and Their Applications (Kōriyama, 1988)*; Ellis Horwood Series Mathematics Applied; Horwood: Chichester, UK, 1988; pp. 329–354.
44. Ismail, M.E.H.; Merkes, E.; Styer, D. A generalization of starlike functions. *Complex Variables Theory Appl.* **1990**, *14*, 77–84. [[CrossRef](#)]
45. Srivastava, H.M. Operators of basic (or q -) calculus and fractional q -calculus and their applications in geometric function theory of complex analysis. *Iran. J. Sci. Technol. Trans. A Sci.* **2020**, *44*, 327–344. [[CrossRef](#)]
46. Jackson, F.H. On q -definite integrals. *Quart. J. Pure Appl. Math.* **1910**, *41*, 193–203.
47. Srivastava, H.M.; Khan, B.; Khan, N.; Ahmad, Q.Z. Coefficient inequalities for q -starlike functions associated with the Janowski functions. *Hokkaido Math. J.* **2019**, *48*, 407–425. [[CrossRef](#)]
48. Srivastava, H.M.; Ahmad, Q.Z.; Khan, N.; Khan, N.; Khan, B. Hankel and Toeplitz determinants for a subclass of q -starlike functions associated with a general conic domain. *Mathematics* **2019**, *7*, 181. [[CrossRef](#)]
49. Srivastava, H.M.; Khan, B.; Khan, N.; Ahmad, Q.Z.; Tahir, M. A generalized conic domain and its applications to certain subclasses of analytic functions. *Rocky Mountain J. Math.* **2019**, *49*, 2325–2346. [[CrossRef](#)]
50. Srivastava, H.M.; Khan, N.; Darus, M.; Rahim, M.T.; Ahmad, Q.Z.; Zeb, Y. Properties of spiral-like close-to-convex functions associated with conic domains. *Mathematics* **2019**, *7*, 706. [[CrossRef](#)]
51. Srivastava, H.M.; Raza, N.; AbuJarad, E.S.A.; Srivastava, G.; AbuJarad, M.H. Fekete-Szegő inequality for classes of (p, q) -starlike and (p, q) -convex functions. *Rev. Real Acad. Cienc. Exactas Fís. Natur. Ser. A Mat. (RACSAM)* **2019**, *113*, 3563–3584. [[CrossRef](#)]
52. Srivastava, H.M.; Tahir, M.; Khan, B.; Ahmad, Q.Z.; Khan, N. Some general classes of q -starlike functions associated with the Janowski functions. *Symmetry* **2019**, *11*, 292. [[CrossRef](#)]
53. Srivastava, H.M.; Tahir, M.; Khan, B.; Ahmad, Q.Z.; Khan, N. Some general families of q -starlike functions associated with the Janowski functions. *Filomat* **2019**, *33*, 2613–2626. [[CrossRef](#)]
54. Srivastava, H.M.; Khan, N.; Khan, S.; Ahmad, Q.Z.; Khan, B. A class of k -symmetric harmonic functions involving a certain q -derivative operator. *Mathematics* **2021**, *9*, 1812. [[CrossRef](#)]
55. Aldawish, I.; Ibrahim, R.W. Solvability of a new q -differential equation related to q -differential inequality of a special type of analytic functions. *Fractal Fract.* **2021**, *5*, 228. [[CrossRef](#)]
56. Zhou, H.; Selvakumaran, K.A.; Sivasubramanian, S.; Purohit, S.D.; Tang, H. Subordination problems for a new class of Bazilevič functions associated with k -symmetric points and fractional q -calculus operators. *AIMS Math.* **2021**, *6*, 8642–8653. [[CrossRef](#)]
57. Ramachandran, C.; Kavitha, D.; Soupramanien, T. Certain bound for q -starlike and q -convex functions with respect to symmetric points. *Int. J. Math. Math. Sci.* **2015**, *2015*, 205682. [[CrossRef](#)]