

12-1-2022

Some Properties of Bazilevič Functions Involving Srivastava–Tomovski Operator

Daniel Breaz

Universitatea 1 Decembrie 1918 din Alba Iulia

Kadhavoor R. Karthikeyan

National University of Science & Technology, Oman

Elangho Umadevi

Zayed University

Alagiriswamy Senguttuvan

National University of Science & Technology, Oman

Follow this and additional works at: <https://zuscholars.zu.ac.ae/works>



Part of the [Mathematics Commons](#)

Recommended Citation

Breaz, Daniel; Karthikeyan, Kadhavoor R.; Umadevi, Elangho; and Senguttuvan, Alagiriswamy, "Some Properties of Bazilevič Functions Involving Srivastava–Tomovski Operator" (2022). *All Works*. 5554. <https://zuscholars.zu.ac.ae/works/5554>

This Article is brought to you for free and open access by ZU Scholars. It has been accepted for inclusion in All Works by an authorized administrator of ZU Scholars. For more information, please contact scholars@zu.ac.ae.

Article

Some Properties of Bazilevič Functions Involving Srivastava–Tomovski Operator

Daniel Breaz ^{1,*}, Kadhavoor R. Karthikeyan ^{2,†}, Elangho Umadevi ^{3,†} and Alagiriswamy Senguttuvan ^{2,†}

¹ Department of Mathematics, “1 Decembrie 1918” University of Alba Iulia, 510009 Alba Iulia, Romania

² Department of Applied Mathematics and Science, National University of Science & Technology, Muscat P.O. Box 620, Oman

³ Department of Mathematics and Statistics, College of Natural and Health Sciences, Zayed University, Abu Dhabi P.O. Box 144534, United Arab Emirates

* Correspondence: dbreaz@uab.ro

† Authors dedicate this article to Professor Hari M. Srivastava on the occasion of his 82nd Birthday.

Abstract: We introduce a new class of Bazilevič functions involving the Srivastava–Tomovski generalization of the Mittag-Leffler function. The family of functions introduced here is superordinated by a conic domain, which is impacted by the Janowski function. We obtain coefficient estimates and subordination conditions for starlikeness and Fekete–Szegő functional for functions belonging to the class.

Keywords: analytic function; Srivastava–Tomovski generalization of Mittag-Leffler function; convex function; Bazilevič function; starlike function; Fekete–Szegő problem; differential subordination



Citation: Breaz, D.; Karthikeyan, K.R.; Umadevi, E.; Senguttuvan, A. Some Properties of Bazilevič Functions Involving Srivastava–Tomovski Operator. *Axioms* **2022**, *11*, 687. <https://doi.org/10.3390/axioms11120687>

Academic Editors: Juan J. Nieto and Sevtap Sümer Eker

Received: 26 October 2022

Accepted: 28 November 2022

Published: 30 November 2022

Publisher’s Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2022 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Researchers have successfully applied the *Mittag-Leffler function and its multi-parameter extensions* to several problems in physics, engineering and other applied sciences. However, the real importance of this function arose from the role it plays in *Fractional Calculus* [1]. The familiar Mittag-Leffler function $E_\alpha(z)$ and its two-parameter version $E_{\alpha,\beta}(z)$ are defined, respectively, by

$$E_\alpha(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\alpha n + 1)} \quad \text{and} \quad E_{\alpha,\beta}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\alpha n + \beta)} \quad (z, \alpha, \beta, \in \mathbb{C}, \operatorname{Re}(\alpha) > 0),$$

where $z \in \Pi = \{z : |z| < 1\}$, \mathbb{C} denotes the sets of complex numbers and $(x)_n$ denotes the Pochhammer symbol defined by

$$(x)_n = \frac{\Gamma(x+n)}{\Gamma(x)} = \begin{cases} 1 & \text{if } n = 0 \\ x(x+1)(x+2) \dots (x+n-1) & \text{if } n \in \mathbb{N}. \end{cases}$$

The Mittag-Leffler function $E_\alpha(z)$ and its two-parameter version $E_{\alpha,\beta}(z)$ were first considered by Gösta Mittag-Leffler in 1903 and A. Wiman in 1905. Refer to Ayub et al. [2], Gorenflo [3], Srivastava [4–6] and Srivastava et al. [7–18] for detailed studies that involve the Mittag-Leffler function.

The Mittag-Leffler function $E_{\alpha,\beta}(z)$ coincides with well-known elementary functions and some special functions. For example,

$$E_{1,2}(z) = \frac{e^z - 1}{z}, \quad E_{3,1}(z) = \frac{1}{2} \left[e^{z^{1/3}} + 2e^{-\frac{1}{2}z^{1/3}} \cos\left(\frac{\sqrt{3}}{2}z^{1/3}\right) \right].$$

Srivastava et al. [13] considered the following family of the multi-index Mittag-Leffler functions as a kernel of some fractional-calculus operators

$$E_{(\alpha_j, \beta_j)_m}^{\gamma, k, \delta, \epsilon}(z) = \sum_{n=0}^{\infty} \frac{(\gamma)_{kn} (\delta)_{\epsilon n}}{\prod_{j=1}^m \Gamma(\alpha_j n + \beta_j)} \frac{z^n}{n!} \tag{1}$$

$$\left(\alpha_j, \beta_j, \gamma, k, \delta, \epsilon \in \mathbb{C}; \operatorname{Re}(\alpha_j) > 0, (j = 1, \dots, m); \operatorname{Re}\left(\sum_{j=1}^m \alpha_j\right) > \operatorname{Re}(k + \epsilon) - 1 \right).$$

Some Higher Transcendental Functions and Related Mittag-Leffler Functions

The well-known Meijer G-function and Fox’s H-function have almost all elementary and special functions as their special cases. Here we will restrict with a brief overview of the Fox–Wright function and Hurwitz–Lerch type zeta functions unification with the Mittag-Leffler function and its multi-parameter extensions.

For $\eta_j \in \mathbb{C} (j = 1, \dots, r)$ and $\nu_j \in \mathbb{C} \setminus Z_0^- = \{0, -1, \dots\} (j = 1, \dots, s)$, the Fox–Wright function ${}_r\Psi_s$, which is defined by (see ([19], Equation (1.6)), ([20], p. 21) and ([21], p. 19))

$${}_r\Psi_s \left[\begin{matrix} (\eta_1, A_1) & \dots & (\eta_r, A_r) \\ (\nu_1, B_1) & \dots & (\nu_s, B_s) \end{matrix} ; z \right] = \sum_{n=0}^{\infty} \frac{\prod_{j=1}^r \Gamma(\eta_j + A_j n)}{\prod_{j=1}^s \Gamma(\nu_j + B_j n)} \frac{z^n}{n!} \tag{2}$$

where $\operatorname{Re}(A_j) > 0, (j = 1, \dots, r)$ and $\operatorname{Re}(B_j) > 0 \in \mathbb{C} (j = 1, \dots, s)$ with $1 + \operatorname{Re}\left(\sum_{j=1}^s B_j - \sum_{j=1}^r A_j\right) \geq 0$. Refer to Srivastava ([5], Definition 2) for a detailed discussion on the convergence of the series (2).

Lin and Srivastava [22] introduced and investigated an interesting generalization of the well-known Hurwitz–Lerch zeta function $\phi(z, s, a)$ in the following form

$$\phi_{\eta, \nu}^{k, \epsilon}(z, m, a) = \sum_{n=0}^{\infty} \frac{(\eta)_{kn}}{(\nu)_{\epsilon n}} \frac{z^n}{(n + a)^m}.$$

In order to derive a direct relationship with the Fox–Wright function, the function $\phi_{\eta, \nu}^{k, \epsilon}(z, m, a)$ was further generalized to (see Srivastava et al. [23])

$$\phi_{\eta_1, \dots, \eta_r; \nu_1, \dots, \nu_s}^{k_1, \dots, k_r; \epsilon_1, \dots, \epsilon_s}(z, m, a) = \sum_{n=0}^{\infty} \frac{\prod_{j=1}^r (\eta_j)_{nk_j}}{n! \prod_{j=1}^s (\nu_j)_{n\epsilon_j}} \frac{z^n}{(n + a)^m}$$

where $\eta_j \in \mathbb{C} (j = 1, \dots, r)$ and $\nu_j \in \mathbb{C} \setminus Z_0^- = \{0, -1, \dots\} (j = 1, \dots, s)$. Refer to [24–26], for a detailed discussion on the convergence.

When $A_j = 1, (j = 1, \dots, r)$ and $B_j = 1, (j = 1, \dots, s)$ in (2), then

$${}_r\Psi_s \left[\begin{matrix} (\eta_1, 1) & \dots & (\eta_r, 1) \\ (\nu_1, 1) & \dots & (\nu_s, 1) \end{matrix} ; z \right] = {}_rF_s \left[\begin{matrix} \eta_1 & \dots & \eta_r \\ \nu_1 & \dots & \nu_s \end{matrix} ; z \right] = \sum_{n=0}^{\infty} \frac{(\eta_1)_n \dots (\eta_r)_n}{(\nu_1)_n \dots (\nu_s)_n} \frac{z^n}{n!}.$$

The function ${}_rF_s$ is the well-known generalized hypergeometric function (see [27,28]). Similarly, we observe from (1) and (2) that

$$E_{(\alpha_j, \beta_j)_m}^{\gamma, k, \delta, \epsilon}(z) = \frac{1}{\Gamma(\gamma)\Gamma(\delta)} {}_2\Psi_m \left[\begin{matrix} (\gamma, k) & (\delta, \epsilon) \\ (\beta_1, \alpha_1) & \dots & (\beta_m, \alpha_m) \end{matrix} ; z \right].$$

A special case of the multi-index Mittag-Leffler function defined by (1), when $m = 2$ corresponding to the Srivastava–Tomovski generalization of the Mittag-Leffler function [29], is given by

$$E_{\alpha, \beta}^{\gamma, k}(z) = \sum_{n=0}^{\infty} \frac{(\gamma)_{nk} z^n}{\Gamma(\alpha n + \beta) n!}, \quad z, \alpha, \beta, \gamma, k \in \mathbb{C}, \operatorname{Re}(\alpha) > 0, \operatorname{Re}(k) > 0,$$

$$= \frac{1}{\Gamma(\gamma)} {}_1\Psi_1 \left[\begin{matrix} (\gamma, k) \\ (\beta, \alpha) \end{matrix}; z \right]. \tag{3}$$

In [29], the authors established that $E_{\alpha, \beta}^{\gamma, k}(z)$ defined by (3) is an entire function in the complex z -plane. The function $E_{\alpha, \beta}^{\gamma, k}(z)$ is called the Srivastava–Tomovski generalization of the Mittag-Leffler function. The function $E_{\alpha, \beta}^{\gamma, 1}(z)$ is popularly known as Prabhakar function or generalized Mittag-Leffler three-parameter function.

In geometric function theory, several researchers have studied the properties of the Srivastava–Tomovski generalization of the Mittag-Leffler function. The most prominent studies pertaining to the Srivastava–Tomovski generalization were by Aouf and Mostafa [30], Attiya [31], Liu [32] and Tomovski et al. [33].

2. Definitions and Preliminaries

Let $\mathcal{H}(d, n)$ be the class of analytic functions having a series of the form $\varphi(z) = d + d_n z^n + d_{n+1} z^{n+1} + \dots$

Let

$$\Lambda_n = \{ \varphi \in \mathcal{H}, \varphi(z) = z + d_{n+1} z^{n+1} + d_{n+2} z^{n+2} + \dots \}$$

and let $\Lambda = \Lambda_1$. For $\varphi \in \Lambda$, Cang and Liu in [34] introduced an operator using the Srivastava–Tomovski generalization of the Mittag-Leffler function ([29]), which, explicitly for $p = 1$, is given by

$$H_{\alpha, \beta}^{\gamma, k} \varphi(z) = z + \sum_{n=2}^{\infty} \frac{\Gamma(\gamma + nk) \Gamma(\alpha + \beta)}{\Gamma(\gamma + k) \Gamma(\alpha n + \beta) n!} d_n z^n. \tag{4}$$

Motivated by [35,36], we now define an operator $\mathfrak{J}_k^m(\alpha, \beta, \gamma) \varphi(z) : \Lambda \rightarrow \Lambda$ is defined by

$$\mathfrak{J}_{k, \lambda}^m(\alpha, \beta, \gamma) \varphi(z) = z + \sum_{n=2}^{\infty} [1 - \lambda + \lambda n]^m \frac{\Gamma(\gamma + nk) \Gamma(\alpha + \beta)}{\Gamma(\gamma + k) \Gamma(\alpha n + \beta) n!} d_n z^n. \tag{5}$$

Remark 1. We note that operator $\mathfrak{J}_{k, \lambda}^m(\alpha, \beta, \gamma) \varphi(z)$ is closely related to the operators studied by Breaz et al. [37], Cang and Liu [34] and Elhaddad et al. [38]. Now here we list some of the special cases:

1. $\mathfrak{J}_{1, \lambda}^m(0, \beta, 1) \varphi(z) = D^m \varphi(z)$, where $D^m \varphi(z)$ is the Al-Oboudi operator (see [39])
2. If we let $\alpha = 0, k = \gamma = 1$ and $\lambda = 1$ in (5), then $\mathfrak{J}_{k, \lambda}^m(\alpha, \beta, \gamma) \varphi(z)$ reduces to the well-known Sălăgean operator.
3. If we let $m = 0$ and $\lambda = 1$ in (5), then $\mathfrak{J}_{k, \lambda}^m(\alpha, \beta, \gamma) \varphi(z)$ reduces to the operator defined and studied by Cang & Liu [34].

Let Υ denote the class of functions having series

$$r(z) = 1 + \sum_{n=1}^{\infty} r_n z^n \quad (z \in \Pi),$$

which satisfies the condition $\operatorname{Re}(r(z)) > 0$. We denote by $\mathcal{S}^*(\eta), \mathcal{C}(\eta)$ and $\mathcal{K}(\eta, \tau)$ ($0 \leq \eta, \tau < 1$) the familiar subclasses of Λ consisting of functions that are, respectively, starlike of order η , convex of order η and close-to-convex of order η and type τ in Π .

Recently, Breaz et al. [40] defined and studied the following function

$$\Gamma(Q, R; p; \sigma; \Psi) = \frac{[(1 + Q)p + \sigma(R - Q)]\Psi(z) + [(1 - Q)p - \sigma(R - Q)]}{[(R + 1)\Psi(z) + (1 - R)],} \tag{6}$$

where $\Psi(z) \in Y$ and has a series representation of the form

$$\Psi(z) = 1 + L_1z + L_2z^2 + \dots \tag{7}$$

A detailed geometric interpretation of $\Gamma(Q, R; p; \sigma; \Psi)$ was discussed by Karthikeyan et al. in [41]. The function $\Gamma(Q, R; p; \sigma; \Psi)$ was mainly motivated by the study of Noor and Malik [42] and Srivastava et al. [43–49].

By making use of the function $\Gamma(Q, R; p; \sigma; \Psi)$, we now define the following.

Definition 1. For $0 \leq \delta < \infty$, a function $\varphi(z) \in \Lambda$ is said to be in $\mathcal{B}_\lambda^m(\alpha, \beta, \gamma; \delta; Q, R; \Psi)$ if and only if for all $\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\chi \in \mathcal{S}^*(0)$ it satisfies the condition

$$\frac{z[\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]'}{[\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]^{1-\delta} [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\chi(z)]^\delta} \prec \Gamma(Q, R; 1; \sigma; \Psi), \quad (z \in \Pi) \tag{8}$$

where $\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\chi(z) \neq 0$ for all $z \in \Pi$.

If we let $m = \sigma = \alpha = 0, k = 1, Q = 1, R = -1$ and $\chi \in \mathcal{S}^*(1 + z/1 - z)$, then $\mathcal{B}_\lambda^m(\alpha, \beta, \gamma; Q, R; \Psi)$ reduces to the class

$$\mathcal{B}(\delta; \Psi) = \left\{ \varphi \in \Lambda : \frac{z\varphi'(z)}{[\varphi(z)]^{1-\delta} [\chi(z)]^\delta} \prec \Psi(z), \quad \chi \in \mathcal{S}^*(0) \right\}.$$

The function $\mathcal{B}(\delta; \Psi)$ was studied by Goyal and Goswami in [50] but with φ and χ belonging to Λ_n .

If we let $\chi(z) = z, \sigma = 0, \Psi(z) = \frac{1+z}{1-z}, Q = 1 - 2\eta$ and $R = -1$ we get the class $\mathcal{B}(\delta, \eta)$

which satisfies the condition $\operatorname{Re} \frac{z^{1-\delta} [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]'}{[\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]^{1-\delta}} > \eta, z \in \Pi$, where $\varphi \in \Lambda$.

Further, on letting $m = \sigma = \alpha = 0$ and $k = 1$ in $\mathcal{B}(\delta, \eta)$, it reduces to the well-known class $\mathcal{B}(\delta)$ (see [51]), which satisfies the condition $\operatorname{Re} \frac{z^{1-\delta}\varphi'(z)}{(\varphi(z))^{1-\delta}} > 0, z \in \Pi$, where $\varphi \in \Lambda$. For recent developments pertaining to the study of Bazilevič functions, refer to [52,53].

Throughout this paper, we let

$$\aleph(1; Q, R; \sigma; \Psi; z) = \frac{[(1 + Q) + \sigma(R - Q)]\Psi(z) + [(1 - Q) - \sigma(R - Q)]}{[(R + 1)\Psi(z) + (1 - R)]}. \tag{9}$$

From ([40], Theorem 2), with

$$w(z) = \frac{1}{2}\rho_1z + \frac{1}{2}\left(\rho_2 - \frac{1}{2}\rho_1^2\right)z^2 + \frac{1}{2}\left(\rho_3 - \rho_1\rho_2 + \frac{1}{4}\rho_1^3\right)z^3 + \dots, \quad z \in \Pi,$$

we can get

$$\begin{aligned} \aleph(1; Q, R; \sigma; w(z)) &= 1 + \frac{L_1\rho_1(Q - R)(1 - \sigma)}{4}z + \\ &\frac{(Q - R)(1 - \sigma)L_1}{4} \left[\rho_2 - \rho_1^2 \left(\frac{(R + 1)L_1 + 2\left(1 - \frac{L_2}{L_1}\right)}{4} \right) \right] z^2 + \dots \end{aligned} \tag{10}$$

Now we will state some results, which we will be using to establish the coefficient inequalities.

Lemma 1 ([54]). Let $\rho(z) = 1 + \sum_{n=1}^{\infty} \rho_n z^n$ be analytic in the unit disc satisfying $\operatorname{Re} \rho(z) > 0$. Then, for each complex number ϑ , we have

$$|\rho_2 - \vartheta \rho_1^2| \leq 2 \max\{1, |2\vartheta - 1|\},$$

the result is sharp for functions given by

$$\rho(z) = \frac{1+z^2}{1-z^2}, \quad \rho(z) = \frac{1+z}{1-z}.$$

Lemma 2 ([55]). If $R(z) = z + \sum_{n=2}^{\infty} c_n z^n \in \mathcal{S}^*$, then $|c_n| \leq n$. Further, for each complex number ϑ we have $|c_3 - \vartheta c_2^2| \leq \max(1, |3 - 4\vartheta|)$ and the result is sharp for the Koebe function

$$r(z) = \frac{z}{(1-z)^2} \quad \text{if} \quad \left| \rho - \frac{3}{4} \right| \geq \frac{1}{4}$$

and for

$$r^{\frac{1}{2}}(z^2) = \frac{z}{1-z^2} \quad \text{if} \quad \left| \rho - \frac{3}{4} \right| \leq \frac{1}{4}.$$

3. Fekete–Szegő Inequalities for the Class $\mathcal{B}_\lambda^m(\alpha, \beta, \gamma; \delta; Q, R; \Psi)$

In this section, we obtain the solution to the Fekete–Szegő problem for functions in class $\mathcal{B}_\lambda^m(\alpha, \beta, \gamma; \delta; Q, R; \Psi)$.

Theorem 1. If $\varphi(z) \in \mathcal{B}_\lambda^m(\alpha, \beta, \gamma; \delta; Q, R; \Psi)$, then we have

$$|d_2| \leq \frac{(Q-R)(1-\sigma)|L_1|}{2|M_2|(1+\delta)} + \frac{2\delta}{(\delta+1)}, \tag{11}$$

and

$$|d_3| \leq \frac{(Q-R)(1-\sigma)|L_1|}{2|M_3|(\delta+2)} \max\{1, |2\Gamma_1 - 1|\} + \frac{\delta}{(\delta+2)} \max\{1, |3 - 4\Gamma_2|\} + \frac{(Q-R)\delta(1-\sigma)|L_1 M_2|}{|M_3|(\delta+1)(\delta+2)}, \tag{12}$$

where Γ_1 and Γ_2 are given by

$$\Gamma_1 = \frac{1}{4} \left[L_1(R+1) + 2 \left(1 - \frac{L_2}{L_1} \right) - \frac{(Q-R)(1-\sigma)L_1(\delta^2 + \delta - 2)}{2(1+\delta)^2} \right]$$

$$\Gamma_2 = \frac{M_2^2}{2M_3} \left[(\delta+1) + \frac{\delta(\delta^2 + \delta - 2)}{(1+\delta)^2} + \frac{2\delta}{(\delta+1)} \right].$$

Further, for all $\vartheta \in \mathbb{C}$ we have

$$\left| d_3 - \vartheta d_2^2 \right| \leq \frac{(Q-R)(1-\sigma)|L_1|}{2|M_3|(\delta+2)} \max\{1, |2\mathcal{H}_1 - 1|\} + \frac{2(Q-R)\delta(1-\sigma)|L_1 M_2|}{(\delta+1)} \left| \frac{1}{2M_3(\delta+2)} - \frac{\vartheta}{M_2(1+\delta)} \right| + \frac{2\delta}{(\delta+2)} \max\{1, |3 - 4\mathcal{H}_2|\}, \tag{13}$$

where \mathcal{H}_1 and \mathcal{H}_2 are given by

$$\mathcal{H}_1 = \frac{1}{4} \left[L_1(R + 1) + 2 \left(1 - \frac{L_2}{L_1} \right) + \left\{ \frac{\vartheta M_3(\delta + 2)}{M_2^2} - \frac{(\delta^2 + \delta - 2)}{2} \right\} \frac{(Q - R)(1 - \sigma)L_1}{(1 + \delta)^2} \right]$$

$$\mathcal{H}_2 = \frac{M_2^2}{2M_3} \left[(\delta + 1) - \frac{2\vartheta\delta(\delta + 2)M_3}{M_2^2(\delta + 1)^2} + \frac{\delta(\delta^2 + \delta - 2)}{(1 + \delta)^2} + \frac{2\delta}{(\delta + 1)} \right].$$

The inequality is sharp for each $\vartheta \in \mathbb{C}$.

Proof. By the definition of $\mathcal{B}_{\lambda}^m(\alpha, \beta, \gamma; \delta; Q, R; \Psi)$, we have

$$\frac{z \left[\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z) \right]'}{\left[\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z) \right]^{1-\delta} \left[\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\chi(z) \right]^\delta} = \aleph(1; Q, R; \sigma; w(z)), \tag{14}$$

where $\aleph(1; Q, R; \sigma; w(z))$ is defined as in (9). The left-hand side of (14) is given by

$$\frac{z \left[\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z) \right]'}{\left[\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z) \right]^{1-\delta} \left[\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\chi(z) \right]^\delta}$$

$$= 1 + [d_2(1 + \delta) - \delta c_2]M_2 z + \left[M_3(\delta + 2) \left(d_3 - \frac{1}{2M_3(\delta + 2)} d_2^2 M_2^2 (\delta^2 + \delta - 2) \right) \right. \\ \left. - d_2 M_2^2 \delta c_2 + M_3 \delta \left(c_3 - \frac{M_2^2}{2M_3} c_2^2 (\delta + 1) \right) \right] z^2 + \dots \tag{15}$$

From (15) and (10), the coefficients of z and z^2 are given by

$$d_2 = \frac{(Q - R)(1 - \sigma)L_1\rho_1}{4M_2(1 + \delta)} + \frac{\delta}{(\delta + 1)}c_2 \tag{16}$$

and

$$d_3 = \frac{(Q - R)(1 - \sigma)L_1}{4M_3(\delta + 2)} \left[\rho_2 - \frac{\rho_1^2}{4} \left(L_1(R + 1) + 2 \left(1 - \frac{L_2}{L_1} \right) - \frac{(Q - R)(1 - \sigma)L_1(\delta^2 + \delta - 2)}{2(1 + \delta)^2} \right) \right] + \frac{c_2(Q - R)\delta(1 - \sigma)L_1\rho_1 M_2}{4M_3(\delta + 1)(\delta + 2)} \\ - \frac{\delta}{(\delta + 2)} \left\{ c_3 - \frac{M_2^2}{2M_3} \left[(\delta + 1) + \frac{\delta(\delta^2 + \delta - 2)}{(1 + \delta)^2} + \frac{2\delta}{(\delta + 1)} \right] c_2^2 \right\}. \tag{17}$$

Using $|\rho_n| \leq 2$ ($n \geq 1$) in (16), we can obtain (11). Using (17) together with Lemma 1, we have

$$|d_3| \leq \frac{(Q - R)(1 - \sigma)|L_1|}{4|M_3|(\delta + 2)} \left| \rho_2 - \frac{\rho_1^2}{4} \left(L_1(R + 1) + 2 \left(1 - \frac{L_2}{L_1} \right) - \frac{(Q - R)(1 - \sigma)L_1(\delta^2 + \delta - 2)}{2(1 + \delta)^2} \right) \right| \\ + \frac{\delta}{(\delta + 2)} \left| c_3 - \frac{M_2^2}{2M_3} \left[(\delta + 1) + \frac{\delta(\delta^2 + \delta - 2)}{(1 + \delta)^2} + \frac{2\delta}{(\delta + 1)} \right] c_2^2 \right| \\ + \frac{|c_2|(Q - R)\delta(1 - \sigma)|L_1\rho_1 M_2|}{4|M_3|(\delta + 1)(\delta + 2)}.$$

Hence the proof of (1).
 To establish (1), we consider

$$\begin{aligned}
 |d_3 - \vartheta d_2^2| &= \left| \frac{(Q-R)(1-\sigma)L_1}{4M_3(\delta+2)} \left[\rho_2 - \frac{\rho_1^2}{4} \left(L_1(R+1) + 2 \left(1 - \frac{L_2}{L_1} \right) \right. \right. \right. \\
 &\quad \left. \left. \left. + \left\{ \frac{\vartheta M_3(\delta+2)}{M_2^2} - \frac{(\delta^2 + \delta - 2)}{2} \right\} \frac{(Q-R)(1-\sigma)L_1}{(1+\delta)^2} \right) \right] \right. \\
 &\quad \left. + \frac{c_2(Q-R)\delta(1-\sigma)L_1\rho_1 M_2}{2(\delta+1)} \left\{ \frac{1}{2M_3(\delta+2)} - \frac{\vartheta}{M_2(1+\delta)} \right\} - \frac{\delta}{(\delta+2)} \{c_3 \right. \right. \\
 &\quad \left. \left. - \frac{M_2^2}{2M_3} \left[(\delta+1) - \frac{2\vartheta\delta(\delta+2)M_3}{M_2^2(\delta+1)^2} + \frac{\delta(\delta^2 + \delta - 2)}{(1+\delta)^2} + \frac{2\delta}{(\delta+1)} \right] c_2^2 \right\} \right|. \tag{18}
 \end{aligned}$$

Applying Lemmas 1 and 2 in (18), we can establish inequality (1). □

We denote by $\mathcal{S}(\eta, \tau)$ ($0 \leq \eta < 1 < \tau$) (see [56]) the class of functions $\varphi \in \Lambda$ satisfying the inequality

$$\eta < \operatorname{Re} \frac{z\varphi'(z)}{\varphi(z)} < \tau, \quad z \in \Pi. \tag{19}$$

Corollary 1 ([57], Theorem 5). *Let $0 \leq \eta < 1 < \tau$ and let the function $\varphi \in \mathcal{S}(\eta, \tau)$. Then, for all $\vartheta \in \mathbb{C}$ we have*

$$\begin{aligned}
 |d_3 - \vartheta d_2^2| &\leq \frac{\tau - \eta}{\pi} \sin \frac{\pi(1-\eta)}{\tau - \eta} \\
 &\cdot \max \left\{ 1; \left| \frac{1}{2} + (1-2\vartheta) \frac{\tau - \eta}{\pi} i + \left(\frac{1}{2} - (1-2\vartheta) \frac{\tau - \eta}{\pi} i \right) e^{2\pi i \frac{1-\eta}{\tau-\eta}} \right| \right\}.
 \end{aligned}$$

Proof. From [56], (19) can be rewritten in the form

$$\frac{z\varphi'(z)}{\varphi(z)} < 1 + \frac{\eta - \tau}{\pi} i \log \left(\frac{1 - e^{2\pi i((1-\eta)/(\eta-\tau))} z}{1 - z} \right) = \mathcal{T}(z).$$

Further, it is known that $\operatorname{Re}\{\mathcal{T}(z)\} > 0$ ($z \in \Pi$) and has a series of the form

$$\mathcal{T}(z) = 1 + \sum_{n=1}^{\infty} L_n z^n, \quad z \in \Pi,$$

where $L_n = \frac{\eta - \tau}{n\pi} i \left(1 - e^{2n\pi i((1-\eta)/(\eta-\tau))} \right)$, $n \geq 1$. Substituting the values of $m = \sigma = \delta = \alpha = 0$, $k = 1$, $Q = 1$, $R = -1$, L_1 and L_2 in Theorem 1 we obtain assertion of our theorem. □

If we take $m = \sigma = \delta = \alpha = 0$, $k = 1$, $Q = 1$ and $R = -1$ in Theorem 1, then we have the following corollary.

Corollary 2 ([58]). *Suppose $\varphi(z) = z + d_2 z^2 + d_3 z^3 + \dots \in \mathcal{S}^*(\Psi)$ ($z \in \Pi$). Then*

$$|d_3 - \vartheta d_2^2| \leq \frac{L_1}{2} \max \left\{ 1; \left| L_1 + \frac{L_2}{L_1} - 2\vartheta L_1 \right| \right\} \quad (\vartheta \in \mathbb{C}).$$

Equality is attained if

$$\varphi(z) = \begin{cases} z \exp \int_0^z [\Psi(t) - 1] \frac{1}{t} dt, & \text{if } \left| L_1 + \frac{L_2}{L_1} - 2\vartheta L_1 \right| \geq 1 \\ z \exp \int_0^z [\Psi(t^2) - 1] \frac{1}{t} dt, & \text{if } \left| L_1 + \frac{L_2}{L_1} - 2\vartheta L_1 \right| \leq 1. \end{cases}$$

4. Subordination Results

In general, we note that $\Gamma(Q, R; 1; \sigma; \Psi)$ need not be convex univalent in Π . However, the function $\Gamma(Q, R; 1; \sigma; \Psi)$ is convex depending on the choice of $\Psi(z)$ (see [41,59]).

Lemma 3. Let ℓ be convex in Π , with $\ell(0) = d, v \neq 0$ and $Re v \geq 0$. If $r \in \mathcal{H}(d, n)$ and

$$r(z) + \frac{zr'(z)}{v} \prec \ell(z),$$

then

$$r(z) \prec q(z) \prec \ell(z),$$

where

$$q(z) = \frac{v}{nz^{v/n}} \int_0^z t^{(v/n)-1} \ell(t) dt.$$

The function q is convex and is the best (d, n) -dominant.

Theorem 2. Let $\varphi, \chi \in \Lambda$ with $\varphi(z), \varphi'(z)$ and $\chi(z) \neq 0$ for all $z \in \Pi \setminus \{0\}$. Further, let $\Gamma(Q, R; 1; \sigma; \Psi)$ be convex univalent in Π with $[\Gamma(Q, R; 1; \sigma; \Psi)]_{z=0} = 1$ and $Re \Gamma(Q, R; 1; \sigma; \Psi) > 0$. Further, suppose that $\chi \in \mathcal{S}^*(0)$ and

$$\left(\frac{z [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]'}{[\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]^{1-\delta} [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\chi(z)]^\delta} \right)^2 \left[3 + 2 \left\{ \frac{z [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]''}{[\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]'} - (1-\delta) \frac{z [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]'}{\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)} - \delta \frac{z [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\chi(z)]'}{\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\chi(z)} \right\} \right] \prec \Gamma(Q, R; 1; \sigma; \Psi). \tag{20}$$

Then

$$\frac{z [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]'}{[\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]^{1-\delta} [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\chi(z)]^\delta} \prec \sqrt{\Omega(z)} \tag{21}$$

where

$$\Omega(z) = \frac{1}{z} \int_0^z \Gamma(Q, R; 1; \sigma; \Psi) dt$$

and K is convex and is the best dominant.

Proof. Let

$$r(z) = \frac{z [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]'}{[\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]^{1-\delta} [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\chi(z)]^\delta} \quad (z \in \Pi; \delta \geq 0),$$

then $r(z) \in \mathcal{H}(1, 1)$ with $r(z) \neq 0$.

By assumption, $\Gamma(Q, R; 1; \sigma; \Psi)$ is convex univalent in Π , which, in turn, implies Ω is convex and univalent in Π . Suppose $T(z) = r^2(z)$, then $T(z) \in \mathcal{H}$ with $T(z) \neq 0$ in Π .

Using logarithmic differentiation, we have

$$\frac{zT'(z)}{T(z)} = 2 \left[1 + \frac{z(\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z))''}{(\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z))'} - (1-\delta) \frac{z(\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z))'}{\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)} - \delta \frac{z(\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\chi(z))'}{\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\chi(z)} \right].$$

Thus by (2), we have

$$T(z) + zT'(z) \prec \ell(z) \quad (z \in \Pi). \tag{22}$$

Now, by Lemma 3, we deduce that

$$T(z) \prec \Omega(z) \prec \ell(z).$$

Since $Re \ell(z) > 0$ and $\Omega(z) \prec \ell(z)$, we also have $Re \Omega(z) > 0$. $\sqrt{\Omega(z)}$ is univalent by virtue of Ω being univalent and $r^2(z) \prec \Omega(z)$ implies that $r(z) \prec \sqrt{\Omega(z)}$, which establishes the assertion. \square

Corollary 3. Let $\varphi, \chi \in \Lambda$ with $\varphi(z), \varphi'(z)$ and $\chi(z) \neq 0$ for all $z \in \Pi \setminus \{0\}$. If $\chi \in \mathcal{S}^*(0)$ and

$$Re \left\{ \left(\frac{z [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]'}{[\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]^{1-\delta} [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\chi(z)]^\delta} \right)^2 \left[2 \frac{z [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]''}{[D_{\lambda,s}^{m,q}(\alpha_1, \beta_1)\varphi(z)]'} - (1-\delta) \frac{z [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]'}{\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)} - \delta \frac{z [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\chi(z)]'}{\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\chi(z)} \right] + 3 \right\} > \xi$$

then

$$Re \left[\frac{z [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]'}{[\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\varphi(z)]^{1-\delta} [\mathfrak{J}_{k,\lambda}^m(\alpha, \beta, \gamma)\chi(z)]^\delta} \right] > \omega(\xi),$$

where $\omega(\xi) = \sqrt{[2(1-\xi) \cdot \log 2 + (2\xi-1)]}$, $(0 \leq \xi < 1)$. The inequality is sharp.

Proof. Let $\sigma = 0, Q = 1, R = -1$ and $\Psi(z) = \frac{1+(2\xi-1)z}{1+z}, 0 \leq \xi < 1$ in Theorem 2, we can easily get the desired result. \square

If we let $m = \delta = \alpha = 0$ and $k = 1$ in Corollary 3, then we have the following

Corollary 4. Let $\varphi \in \Lambda$ with $\varphi'(z)$ and $\varphi(z) \neq 0$ for all $z \in \Pi \setminus \{0\}$. If

$$Re \left\{ \left(\frac{z\varphi'(z)}{\varphi(z)} \right)^2 \left[3 + \frac{2z\varphi''(z)}{\varphi'(z)} - \frac{2z\varphi'(z)}{\varphi(z)} \right] \right\} > \xi,$$

then

$$Re \frac{z\varphi'(z)}{\varphi(z)} > \omega(\xi),$$

where $\omega(\xi) = \sqrt{[2(1-\xi) \cdot \log 2 + (2\xi-1)]}$, $(0 \leq \xi < 1)$. This inequality is sharp.

If we let $\delta = 1, m = \alpha = 0$ and $k = 1$ in Corollary 3, then we have the following

Corollary 5. If $\varphi, \chi \in \Lambda$ satisfies

$$Re \left\{ \left(\frac{z\varphi'(z)}{\chi(z)} \right)^2 \left[3 + \frac{2z\varphi''(z)}{\varphi'(z)} - \frac{2z\chi'(z)}{\chi(z)} \right] \right\} > \xi,$$

with $\chi \in \mathcal{S}^*(0)$ and $\chi(z) \neq 0$ for all $z \in \Pi \setminus \{0\}$, then

$$\operatorname{Re} \frac{z\varphi'(z)}{\chi(z)} > \omega(\xi),$$

where $\omega(\xi) = \sqrt{[2(1-\xi) \cdot \log 2 + (2\xi-1)]}$, ($0 \leq \xi < 1$). The inequality is sharp.

5. Conclusions

The main purpose of this present study is to obtain the coefficient inequality for the class of Bazilevič functions, which is computationally cumbersome. To add more versatility to our study, we have studied a class of Bazilevič functions involving the Mittag-Leffler functions. Coefficient inequality, solutions to the Fekete–Szegő problem and sufficient conditions for starlikeness are the primary results of this paper. We have pointed out appropriate connections that we investigated here, together with those in several interconnected earlier works.

We note that this study can be extended by taking a trigonometric function, exponential function, Legendre polynomial, Chebyshev polynomial, Fibonacci sequence or q -Hermite polynomial instead of considering $\psi(z)$ as in (7).

Author Contributions: Conceptualization, D.B., K.R.K., E.U. and A.S.; methodology, D.B., K.R.K., E.U. and A.S.; software, D.B., K.R.K., E.U. and A.S.; validation, D.B., K.R.K., E.U. and A.S.; formal analysis, D.B., K.R.K., E.U. and A.S.; investigation, D.B., K.R.K., E.U. and A.S.; resources, D.B., K.R.K., E.U. and A.S.; data curation, D.B., K.R.K., E.U. and A.S.; writing—original draft preparation, D.B., K.R.K., E.U. and A.S.; writing—review and editing, D.B., K.R.K., E.U. and A.S.; visualization, D.B., K.R.K., E.U. and A.S.; supervision, D.B., K.R.K., E.U. and A.S.; project administration, D.B., K.R.K., E.U. and A.S. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Not applicable.

Acknowledgments: The authors would like to thank all the reviewers for their helpful comments and suggestions, which helped us remove the mistakes and also led to improvement in the presentation of the results.

Conflicts of Interest: The authors declare that they have no conflict of interest.

References

1. Kilbas, A.A.; Srivastava, H.M.; Trujillo, J.J. *Theory and Applications of Fractional Differential Equations*; North-Holland Mathematics Studies, 204; Elsevier Science B.V.: Amsterdam, The Netherlands, 2006.
2. Ayub, U.; Mubeen, S.; Abdeljawad, T.; Rahman, G.; Nisar, K.S. The new Mittag-Leffler function and its applications. *J. Math.* **2020**, *2020*, 2463782. [[CrossRef](#)]
3. Gorenflo, R.; Kilbas, A.A.; Rogosin, S.V. On the generalized Mittag-Leffler type functions. *Integral Transform. Spec. Funct.* **1998**, *7*, 215–224.
4. Srivastava, H.M. A survey of some recent developments on higher transcendental functions of analytic number theory and applied mathematics. *Symmetry* **2021**, *13*, 2294. [[CrossRef](#)]
5. Srivastava, H.M. An introductory overview of fractional-calculus operators based upon the Fox-Wright and related higher transcendental functions. *J. Adv. Engrg. Comput.* **2021**, *5*, 135–166. [[CrossRef](#)]
6. Srivastava, H.M. Some parametric and argument variations of the operators of fractional calculus and related special functions and integral transformations. *J. Nonlinear Convex Anal.* **2021**, *22*, 1501–1520.
7. Srivastava, H.M.; Kumar, A.; Das, S.; Mehrez, K. Geometric properties of a certain class of Mittag-Leffler-type functions. *Fractal Fract.* **2022**, *6*, 54. [[CrossRef](#)]
8. Srivastava, H.M.; Alomari, A.-K.N.; Saad, K.M.; Hamanah, W.M. Some Dynamical models involving fractional-order derivatives with the Mittag-Leffler type kernels and their applications based upon the Legendre spectral collocation method. *Fractal Fract.* **2021**, *5*, 131. [[CrossRef](#)]

9. Srivastava, H.M.; El-Deeb, S.M. Fuzzy differential subordinations based upon the Mittag-Leffler type Borel distribution. *Symmetry* **2021**, *13*, 1023. [[CrossRef](#)]
10. Srivastava, H.M.; Kashuri, A.; Mohammed, P.O.; Alsharif, A.M.; Guirao, J.L.G. New Chebyshev type inequalities via a general family of fractional integral operators with a modified Mittag-Leffler kernel. *AIMS Math.* **2021**, *6*, 11167–11186. [[CrossRef](#)]
11. Srivastava, H.M.; Murugusundaramoorthy, G.; El-Deeb, S.M. Faber polynomial coefficient estimates of bi-close-to-convex functions connected with the Borel distribution of the Mittag-Leffler type. *J. Nonlinear Var. Anal.* **2021**, *5*, 103–118.
12. Srivastava, H.M.; Fernandez, A.; Baleanu, D. Some new fractional-calculus connections between Mittag-Leffler functions. *Mathematics* **2019**, *7*, 485. [[CrossRef](#)]
13. Srivastava, H.M.; Bansal, M.; Harjule, P. A study of fractional integral operators involving a certain generalized multi-index Mittag-Leffler function. *Math. Methods Appl. Sci.* **2018**, *41*, 6108–6121. [[CrossRef](#)]
14. Srivastava, H.M.; Kilicman, A.; Zainab, A.E.; Ibrahim, A.E. Generalized convolution properties based on the modified Mittag-Leffler function. *J. Nonlinear Sci. Appl.* **2017**, *10*, 4284–4294. [[CrossRef](#)]
15. Srivastava, H.M.; Aliev, N.A.; Mammadova, G.H.; Aliev, F. A Some remarks on the paper, entitled “Fractional and operational calculus with generalized fractional derivative operators and Mittag-Leffler type functions. *TWMS J. Pure Appl. Math.* **2017**, *8*, 112–114.
16. Srivastava, H.M.; Frasin, B.A.; Pescar, V. Univalence of integral operators involving Mittag-Leffler functions. *Appl. Math. Inf. Sci.* **2017**, *11*, 635–641 [[CrossRef](#)]
17. Srivastava, H.M. Some families of Mittag-Leffler type functions and associated operators of fractional calculus (survey). *TWMS J. Pure Appl. Math.* **2016**, *7*, 123–145.
18. Srivastava, H.M. On an extension of the Mittag-Leffler function. *Yokohama Math. J.* **1968**, *16*, 77–88.
19. Srivastava, H.M. Some Fox-Wright generalized hypergeometric functions and associated families of convolution operators. *Appl. Anal. Discret. Math.* **2007**, *1*, 56–71.
20. Srivastava, H.M.; Karlsson, P.W. *Multiple Gaussian Hypergeometric Series*; Ellis Horwood Series: Mathematics and its Applications; Ellis Horwood Ltd.: Chichester, UK, 1985.
21. Srivastava, H.M.; Gupta, K.C.; Goyal, S.P. *The H-Functions of One and Two Variables*; South Asian Publishers Pvt. Ltd.: New Delhi, India, 1982.
22. Lin, S.-D.; Srivastava, H.M. Some families of the Hurwitz-Lerch Zeta functions and associated fractional derivative and other integral representations. *Appl. Math. Comput.* **2004**, *154*, 725–733. [[CrossRef](#)]
23. Srivastava, H.M.; Saxena, R.K.; Pogány, T.K.; Saxena, R. Integral and computational representations of the extended Hurwitz-Lerch zeta function. *Integral Transform. Spec. Funct.* **2011**, *22*, 487–506. [[CrossRef](#)]
24. Srivastava, H.M. Generating relations and other results associated with some families of the extended Hurwitz-Lerch Zeta functions. *SpringerPlus* **2013**, *2*, 67.
25. Srivastava, H.M.; Jankov, D.; Pogány, T.K.; Saxena, R.K. Two-sided inequalities for the extended Hurwitz-Lerch zeta function. *Comput. Math. Appl.* **2011**, *62*, 516–522. [[CrossRef](#)]
26. Răducanu, D.; Srivastava, H.M. A new class of analytic functions defined by means of a convolution operator involving the Hurwitz-Lerch zeta function. *Integral Transform. Spec. Funct.* **2007**, *18*, 933–943. [[CrossRef](#)]
27. Dziok, J.; Srivastava, H.M. Classes of analytic functions associated with the generalized hypergeometric function. *Appl. Math. Comput.* **1999**, *103*, 1–13. [[CrossRef](#)]
28. Dziok, J.; Srivastava, H.M. Certain subclasses of analytic functions associated with the generalized hypergeometric function. *Integral Transform. Spec. Funct.* **2003**, *14*, 7–18.
29. Srivastava, H.M.; Tomovski, Ž. Fractional calculus with an integral operator containing a generalized Mittag-Leffler function in the kernel. *Appl. Math. Comput.* **2009**, *211*, 198–210. [[CrossRef](#)]
30. Aouf, M.K.; Mostafa, A.O. Certain inequalities of meromorphic univalent functions associated with the Mittag-Leffler function. *J. Appl. Anal.* **2019**, *25*, 173–178. [[CrossRef](#)]
31. Attiya, A.A. Some applications of Mittag-Leffler function in the unit disk. *Filomat* **2016**, *30*, 2075–2081. [[CrossRef](#)]
32. Liu, J.-L. New applications of the Srivastava–Tomovski generalization of the Mittag-Leffler function. *Iran. J. Sci. Technol. Trans. A Sci.* **2019**, *43*, 519–524. [[CrossRef](#)]
33. Tomovski, Ž.; Hilfer, R.; Srivastava, H.M. Fractional and operational calculus with generalized fractional derivative operators and Mittag-Leffler type functions. *Integral Transform. Spec. Funct.* **2010**, *21*, 797–814. [[CrossRef](#)]
34. Cang, Y.-L.; Liu, J.-L. A family of multivalent analytic functions associated with Srivastava–Tomovski generalization of the Mittag-Leffler function. *Filomat* **2018**, *32*, 4619–4625. [[CrossRef](#)]
35. Karthikeyan, K.R.; Reddy, K.A.; Murugusundaramoorthy, G. On classes of Janowski functions associated with a conic domain. *Ital. J. Pure Appl. Math.* **2022**, *47*, 684–698.
36. Reddy, K.A.; Karthikeyan, K.R.; Murugusundaramoorthy, G. Inequalities for the Taylor coefficients of spirallike functions involving q -differential operator. *Eur. J. Pure Appl. Math.* **2019**, *12*, 846–856. [[CrossRef](#)]
37. Breaz, D.; Karthikeyan, K.R.; Umadevi, E. Subclasses of Multivalent Meromorphic functions with a pole of order p at the origin. *Mathematics* **2022**, *10*, 600. [[CrossRef](#)]
38. Elhaddad, S.; Aldweby, H.; Darus, M. On certain subclasses of analytic functions involving differential operator. *Jnānābha* **2018**, *48*, 55–64.

39. Al-Oboudi, F.M. On univalent functions defined by a generalized Sălăgean operator. *Int. J. Math. Math. Sci.* **2004**, *27*, 1429–1436. [[CrossRef](#)]
40. Breaz, D.; Karthikeyan, K.R.; Senguttuvan, A. Multivalent prestarlike functions with respect to symmetric points. *Symmetry* **2022**, *14*, 20. [[CrossRef](#)]
41. Karthikeyan, K.R.; Lakshmi, S.; Varadharajan, S.; Mohankumar, D.; Umadevi, E. Starlike functions of complex order with respect to symmetric points defined using higher order derivatives. *Fractal Fract.* **2022**, *6*, 116. [[CrossRef](#)]
42. Noor, K.I.; Malik, S.N. On coefficient inequalities of functions associated with conic domains. *Comput. Math. Appl.* **2011**, *62*, 2209–2217. [[CrossRef](#)]
43. Srivastava, H.M.; Ahmad, Q.Z.; Khan, N.; Khan, N.; Khan, B. Hankel and Toeplitz determinants for a subclass of q -starlike functions associated with a general conic domain. *Mathematics* **2019**, *7*, 181. [[CrossRef](#)]
44. Srivastava, H.M.; Khan, B.; Khan, N.; Ahmad, Q.Z. Coefficient inequalities for q -starlike functions associated with the Janowski functions. *Hokkaido Math. J.* **2019**, *48*, 407–425. [[CrossRef](#)]
45. Srivastava, H.M.; Khan, B.; Khan, N.; Ahmad, Q.Z.; Tahir, M. A generalized conic domain and its applications to certain subclasses of analytic functions. *Rocky Mt. J. Math.* **2019**, *49*, 2325–2346. [[CrossRef](#)]
46. Srivastava, H.M.; Khan, N.; Darus, M.; Rahim, M.T.; Ahmad, Q.Z.; Zeb, Y. Properties of spiral-like close-to-convex functions associated with conic domains. *Mathematics* **2019**, *7*, 706. [[CrossRef](#)]
47. Srivastava, H.M.; Raza, N.; AbuJarad, E.S.A.; Srivastava, G.; AbuJarad, M.H. Fekete-Szegö inequality for classes of (p, q) -starlike and (p, q) -convex functions. *Rev. Real Acad. Cienc. Exactas Fís. Natur. Ser. A Mat. (RACSAM)* **2019**, *113*, 3563–3584. [[CrossRef](#)]
48. Srivastava, H.M.; Tahir, M.; Khan, B.; Ahmad, Q.Z.; Khan, N. Some general classes of q -Starlike functions associated with the Janowski functions. *Symmetry* **2019**, *11*, 292. [[CrossRef](#)]
49. Srivastava, H.M.; Tahir, M.; Khan, B.; Ahmad, Q.Z.; Khan, N. Some general families of q -starlike functions associated with the Janowski functions. *Filomat* **2019**, *33*, 2613–2626. [[CrossRef](#)]
50. Goyal, S.P.; Goswami, P. On sufficient conditions for analytic functions to be Bazilevič. *Complex Var. Elliptic Equ.* **2009**, *54*, 485–492. [[CrossRef](#)]
51. Bazilevič, I.E. On a case of integrability in quadratures of the Loewner-Kufarev equation. *Mat. Sb.* **1955**, *37*, 471–476.
52. Srivastava, H.M.; Wanas, A.K.; Güney, H.Ö. New families of bi-univalent functions associated with the Bazilevič functions and the λ -pseudo-starlike functions. *Iran. J. Sci. Technol. Trans. A Sci.* **2021**, *45*, 1799–1804. [[CrossRef](#)]
53. Wanas, A.K.; Srivastava, H.M. Differential sandwich theorems for Bazilevič function defined by convolution structure. *Turk. J. Ineq.* **2020**, *4*, 10–21.
54. Ma, W.C.; Minda, D. A unified treatment of some special classes of univalent functions. In *Conf. Proc. Lecture Notes Anal., Proceedings of the Conference on Complex Analysis, Tianjin, China, 19–23 June 1992*; International Press: Cambridge, MA, USA, 1992; pp. 157–169.
55. Koepf, W. On the Fekete-Szegö problem for close-to-convex functions. *Proc. Am. Math. Soc.* **1987**, *101*, 89–95.
56. Kuroki, K.; Owa, S. Notes on new class for certain analytic functions. *RIMS Kokyuroku* **2011**, *1772*, 21–25.
57. Sim, Y.J.; Kwon, O.S. Notes on analytic functions with a bounded positive real part. *J. Inequal. Appl.* **2013**, *2013*, 370. [[CrossRef](#)]
58. Tu, Z.; Xiong, L. Unified solution of Fekete-Szegö problem for subclasses of starlike mappings in several complex variables *Math. Slovaca* **2019**, *69*, 843–856. [[CrossRef](#)]
59. Breaz, D.; Karthikeyan, K.R.; Murugusundaramoorthy, G. Bazilevič functions of complex order with respect to symmetric points. *Fractal Fract.* **2022**, *6*, 316. [[CrossRef](#)]